

## Systematic Review

# Is Unilateral Percutaneous Kyphoplasty Superior to Bilateral Percutaneous Kyphoplasty for Osteoporotic Vertebral Compression Fractures? Evidence from a Systematic Review of Discordant Meta-Analyses

Xiaofeng Chen, MD<sup>1</sup>, Weijun Guo, MD<sup>2</sup>, Qian Li, MD<sup>1</sup>, Zhicong Ou, MD<sup>1</sup>, Zehui Lao, MD<sup>1</sup>, Yi Liu, MD<sup>1</sup>, Chenghua Zhu, MD<sup>1</sup>, Zhuangxun Han, MD<sup>1</sup>, Xueyuan Chu, MD<sup>1</sup>, and Dongling Cai, MD<sup>1</sup>

From: <sup>1</sup>Department of Orthopedics, Panyu Hospital of Chinese Medicine, China; <sup>2</sup>Guangzhou University of Chinese Medicine, China

Address Correspondence: Dongling Cai, MD  
Department of Orthopedics, Panyu Hospital of Chinese Medicine, No. 65, Qiaodong Road, Panyu District, Guangzhou 511400, China  
E-mail: panyucaidongling@126.com

Disclaimer: There was no external funding in the preparation of this manuscript. Conflict of interest: Each author certifies that he or she, or a member of his or her immediate family, has no commercial association (i.e., consultancies, stock ownership, equity interest, patent/licensing arrangements, etc.) that might pose a conflict of interest in connection with the submitted manuscript.

Manuscript received: 07-21-2017  
Revised manuscript received: 11-23-2017  
Accepted for publication: 02-01-2018

Free full manuscript: [www.painphysicianjournal.com](http://www.painphysicianjournal.com)

**Background:** Several meta-analyses have been performed to compare unilateral percutaneous kyphoplasty (PKP) and bilateral PKP in the treatment of osteoporotic vertebral compression fractures (OVCFs), but inconsistencies in the results have led to questions as to which technique is preferable.

**Objective:** This study was designed to clarify the benefits and disadvantages of unilateral PKP versus bilateral PKP as found in numerous discordant meta-analyses and thereby present surgical treatment recommendations for OVCFs considering the current best evidence.

**Study Design:** Systematic review/Meta-analysis.

**Methods:** Meta-analyses on unilateral and bilateral PKP for OVCFs were included by searching Pubmed, Embase, and Cochrane library. Meta-analysis quality was assessed using Oxford Levels of Evidence and Assessment of Multiple Systematic Reviews (AMSTAR). The Jadad decision algorithm was used to identify the best evidence.

**Results:** Eight eligible meta-analyses were included, 7 of which were Level-II evidence and one was Level-III evidence. The AMSTAR scores varied from 7 to 8. The Jadad decision algorithm suggested that the best meta-analysis should be selected depending upon publication characteristics and methodology of primary studies, language restrictions, and whether data analysis was performed on individual patients. The best available evidence indicated that both unilateral and bilateral PKP could receive similar good clinical and radiological outcomes. However, without increasing the risk of complications, unilateral PKP required shorter surgical time and less cement volume, offering better pain relief and quality of life at post-operative short term follow-ups.

**Limitations:** Primary studies had defects in their methodologies.

**Conclusions:** Unilateral PKP appears to be superior to bilateral PKP in the treatment of OVCFs.

**Key words:** Osteoporotic vertebral compression fractures, percutaneous kyphoplasty, meta-analysis

**Pain Physician 2018; 21:327-336**

**O**steoporotic vertebral compression fractures (OVCFs) are common fractures in the elderly, and affect approximately 1.4 million patients, with a predominance in women (1). These

fractures frequently lead to decreased mobility and quality of life (2,3). OVCFs have traditionally been managed by conservative treatment such as bed rest. About two-thirds of these patients improve

(2). However, complications from long-term bed-rest can worsen patients' conditions. Moreover, severe pain and progressive deformity are usually reported in patients treated by conservative treatment (2,4). Thus, percutaneous kyphoplasty (PKP) as a safe and effective alternative has been widely performed for OVCFs.

The standard procedure for PKP used a bipedicular approach with two balloon tamps (2,5). PKP using a unipedicular approach has been increasingly performed over the recent years due to reported outcomes comparable to bilateral PKP (6-12). Several randomized clinical trials (RCTs) have been done comparing the effectiveness and safety between unilateral and bilateral PKP, but results have been conflicting (2,13-32). The optimal percutaneous technique for OVCFs remains controversial. Meta-analyses have also been conducted to compare and contrast these 2 techniques. Likewise, results from these meta-analyses have also been conflicting (5-12). Although several studies have reported no differences in treatment effectiveness between unilateral and bilateral PKP in the treatment of OVCFs (7,10,12), other studies have concluded that unilateral PKP is superior to bilateral PKP (5,6,8,9,11). Such conflicting results have created controversy for decision makers (e.g., policy-makers, doctors, patients) who depend on high quality evidence when deciding upon an optimal OVCFs treatment.

In recent years, there has been an increasing number of systematic reviews on discrepant and overlapping meta-analyses (33-38). However, to our knowledge, none of these studies have compared the effectiveness between unilateral and bilateral PKP in the treatment of OVCFs. We conducted such a systematic review in an effort to provide operative recommendations based on the best available evidence.

## **METHODS**

This study was performed following the Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) statement (39) and previous similar reports (40-43).

### **Study Retrieval**

A computer search was done using PubMed, Embase, and the Cochrane Library on January 12, 2017. The following key words were used: kyphoplasty, systematic review, and meta-analysis. Two investigators individually performed the computer search. The reference lists of the meta-analyses were used to identify relevant

studies. The titles and abstracts were checked for potential eligible studies. The full versions were obtained when the data was insufficient. Any disagreement was resolved by a third investigator.

### **Selection Criteria**

The inclusion criteria were as follows: (1) meta-analyses comparing unilateral PKP with bilateral PKP for OVCFs; (2) articles published in English; (3) comparison of one or more outcomes, such as functional outcomes and complications. Abstracts of meetings, letters to the editor, correspondence, and systematic reviews without meta-analysis were excluded.

### **Information Extraction**

The basic information was independently extracted by 2 investigators: the first author, publication year, database searched, the design of primary trial, the number of included studies and RCTs, I2, and outcomes. Disagreement between the investigators was eliminated by discussion with a third investigator.

### **Quality Assessment**

Two investigators were assigned to evaluate the methodological quality of meta-analysis using Oxford Levels of Evidence (44) and the Assessment of Multiple Systematic Reviews (AMSTAR) tool (45). AMSTAR is thought to be a measurement instrument of study methodology, with reliability, validity, and responsibility (46,47). Any disagreement between the investigators was settled based on discussion and, when necessary, with arbitration by a third investigator.

### **Implementation of Jadad Decision Algorithm**

Jadad et al (48) summarized the sources of disagreement among systematic reviews: comprising clinical questions, inclusion and exclusion criteria, abstracting data, quality evaluation, data synthesis, and statistical analysis. This decision algorithm guides the researchers to find treatment recommendations among discordant meta-analyses (40-43). Three investigators individually assessed the meta-analyses by this algorithm and came to an agreement on which meta-analysis generated the current best evidence.

## **RESULTS**

### **Study Retrieval**

The search strategy found a total of 217 records from the databases. The selection process is shown in

Fig. 1. As a result, 8 meta-analyses were included (5-12). The general information is summarized in Table 1. Three studies were published in 2013 (8,10,11), 2 studies were published in 2014 (9,12), one study was published in 2015 (5), and the other 2 studies were published in 2016 (6,7). The meta-analyses included 3 to 14 primary studies (Table 2).

### Search Methodology

Three meta-analyses restricted the publication language to English (5,6,9), while the others reported no language limitation (7,8,10-12). A Cochrane Library search was obtained by all included meta-analyses. However, the use of PubMed, Medline, Embase, OVID, and Web of Science was discordant (Table 3).




Table 1. Characteristics of the included meta-analyses.

First author	Date of publication	Journal	Date of last literature search	No. of included studies	No. of included RCTs
Li LH (8)	2013	Chin Med J (Engl)	March, 2013	7	7
Lin J (11)	2013	Pain Physician	July, 2012	3	3
Yang LY (10)	2013	Pain Physician	September, 2012	4	4
Chen H (12)	2014	Orthopedics	April, 2013	14	14
Huang Z (9)	2014	Clin Orthop Relat Res	June, 2013	5	5
Feng H (5)	2015	J Orthop Res	January, 2015	12	12
Cheng X (7)	2016	Eur Spine J	April, 2015	8	5
Sun H (6)	2016	Pain Physician	June, 2015	6	6

Table 2. Primary studies included in meta-analyses.

First author (Year)	Li LH (2013)	Lin J (2013)	Yang LY (2013)	Chen H (2014)	Huang Z (2014)	Feng H (2015)	Cheng X (2016)	Sun H (2016)
Chung HJ (2008) (15)	+	+	+	+	+	+	+	+
Gu XH (2009) (22)	+			+				
Chen C (2010) (25)	+		+	+	+	+	+	+
Jiang Y (2010) (24)				+				
Chen L (2011) (28)	+	+	+	+	+	+	+	+
Chen C (2011) (16)	+	+	+	+	+	+	+	+
Zhang B (2012) (30)				+				
Luo XL (2012) (32)	+			+				
Li Q (2012) (31)				+				
Li GZ (2012) (14)				+				
Feng J (2012) (13)				+				
YX H (2012) (19)				+				
Feng J (2013) (18)						+		
Mao JG (2013) (29)						+		
Zhai HL (2013) (20)						+		
Huang S (2013) (17)				+				
Rebolledo BJ (2013) (21)	+			+	+	+	+	+
He CJ (2014) (23)						+		
Lin XL (2014) (27)						+		
XJ L (2014) (26)						+		
Yan L (2014) (2)						+		+

Table 3. Search methodology of the included studies.

First author (year)	Restriction of publication language	Restriction of publication status	Search database						
			PubMed	Medline	Embase	Cochrane Library	OVID	Web of Science	Others
Li LH (2013) (8)	No	Yes	+			+			+
Lin J (2013) (11)	No	NR	+	+	+	+			+
Yang LY (2013) (10)	No	NR		+	+	+	+		+
Chen H (2014) (12)	No	NR	+		+	+		+	+
Huang Z (2014) (9)	Yes	Yes	+	+	+	+		+	+
Feng H (2015) (5)	Yes	Yes	+	+	+	+			+
Cheng X (2016) (7)	No	Yes	+	+	+	+	+	+	+
Sun H (2016) (6)	Yes	Yes	+	+	+	+		+	+

Table 4. Methodological information for the included studies.

First author (year)	Design of included studies	Level of evidence	Software	GRADE use	Subgroup analysis	Sensitivity analysis
Li LH (2013) (8)	RCT or quasi-RCT	Level II	Stata	No	No	Yes
Lin J (2013) (11)	RCT	Level II	RevMan	No	No	No
Yang LY (2013) (10)	RCT	Level II	RevMan	No	No	Yes
Chen H (2014) (12)	RCT	Level II	RevMan	Yes	No	No
Huang Z (2014) (9)	RCT or PCS	Level II	RevMan	Yes	No	Yes
Feng H (2015) (5)	RCT	Level II	RevMan	No	No	No
Cheng X (2016) (7)	RCT or non-RCT	Level III	RevMan	No	No	No
Sun H (2016) (6)	RCT	Level II	RevMan	Yes	No	Yes

RCT, Randomized clinical trial; PCS, Prospective comparative study

Table 5. AMSTAR scores for the included studies.

Items	Li LH (2013) (8)	Lin J (2013) (11)	Yang LY (2013) (10)	Chen H (2014) (12)	Huang Z (2014) (9)	Feng H (2015) (5)	Cheng X (2016) (7)	Sun H (2016) (6)
1. Was an a priori design provided?	0	0	0	0	0	0	0	0
2. Was there duplicate study selection and data extraction?	1	1	1	1	1	1	1	1
3. Was a comprehensive literature search performed?	1	1	1	1	1	1	1	1
4. Was the status of publication (i.e., grey literature) used as an inclusion criterion?	0	0	0	0	0	0	0	0
5. Was a list of studies (included and excluded) provided?	0	1	1	0	0	0	0	1
6. Were the characteristics of the included studies provided?	1	1	1	1	1	1	1	1
7. Was the scientific quality of the included studies assessed and documented?	1	1	1	1	1	1	1	1
8. Was the scientific quality of the included studies used appropriately in formulating conclusions?	0	0	0	0	1	1	1	1
9. Were the methods used to combine the findings of studies appropriate?	1	1	1	1	1	1	1	1
10. Was the likelihood of publication bias assessed?	1	0	1	1	0	1	0	0
11. Was the conflict of interest stated?	1	1	1	1	1	1	1	1
Total scores	7	7	8	7	7	8	7	8

### Study Quality

In light of Oxford Levels of Evidence, seven studies were evaluated as Level-II (5,6,8-12), while one study was regarded as Level-III (Table 4) (7). The GRADE was applied in 3 studies (6,9,12). The AMSTAR scores ranged from 7 to 8 (Table 5).

### Heterogeneity Evaluation

Heterogeneity was evaluated by statistical method using the I<sup>2</sup> statistic value (Table 6). Four meta-analyses performed the subgroup analyses (Table 4) (6,8-10). The I<sup>2</sup> values are shown in Table 6. The heterogeneities are acceptable for most results.

Table 6.  $I^2$  statistic value of each variable in each meta-analysis.

Items	Li LH (2013) (8)	Lin J (2013) (11)	Yang LY (2013) (10)	Chen H (2014) (12)	Huang Z (2014) (9)	Feng H (2015) (5)	Cheng X (2016) (7)	Sun H (2016) (6)
Short-term VAS	0%	0%	0%	0%	0%	0%	0%	55%
Middle-term VAS				0%		0%		
Long-term VAS	0%	0%	0%	0%	0%	0%	0%	13%
Short-term ODI	NR			72%	NR	36%	31%	
Middle-term ODI								0%
Long-term ODI	NR			91%	NR			0%
Short-term SF-36 general health benefit						0%		0%
Short-term SF-36 bodily pain relief						9%		
Long-term SF-36						0%		
Surgery time	95%	17%	0%	32%	0%	14%	95%	77%
Cement volume	50%	0%		38%		44%	96%	0%
Cement leakage	40%	45%	28%	27%	41%	0%	23%	22%
Kyphosis angle reduction	0%			42%	85%	85%		93%
Restoration rate	94%					94%	0%	
Short-term anterior vertebral height restoration						0%		
Long-term anterior vertebral height restoration				0%	91%	18%		88%
Middle vertebral height restoration				25%		0%		
Cobb's angle recovery						50%		
X-ray exposure frequency							98%	
Vertebral height loss			68%			0%		68%
Adjacent vertebral fracture			34%		39%	0%	34%	0%

VAS, Visual analog scale; ODI, Oswestry disability index

### Results of Jadad Decision Algorithm

The outcomes of each meta-analysis are listed in Fig. 2. Given that the meta-analyses studied the same question, but did not use the same references or eligibility criteria, the Jadad decision algorithm suggested that the best available evidence should be chosen based upon the publication characteristics of the primary studies, the methodology of the primary studies, the language restrictions, and whether data for analysis were obtained from individual patients (Fig. 3). Hence, a high-quality meta-analysis reported by Feng et al (5) was eventually selected. This study found that unilateral PKP significantly decreased surgery time and cement volume and had a better result of pain relief (visual analog scale (VAS)) as well as quality of life (bodily pain relief and general health benefit of SF-36)

at post-operative short-term follow-ups. Bilateral PKP had a better restoration of anterior vertebral height than unilateral PKP in short-term follow-up, while no difference was observed in long-term follow-up. The differences between unilateral and bilateral PKP were not significant in middle- and long-term VAS, short-term Oswestry disability index (ODI), long-term SF-36, cement leakage, long-term kyphosis angle reduction, restoration rate, long-term anterior vertebral height restoration, middle vertebral height restoration, Cobb's angle recovery, vertebral height loss, and adjacent vertebral fracture.

### DISCUSSION

OVCFs can be managed by nonsurgical or surgical procedure (4). Surgical treatments, including unilateral




Fig. 2. Results of the included meta-analyses.


PKP and bilateral PKP, are widely performed in clinical settings because poor outcomes are reported in patients with conservative treatment (2,13-32). Unilateral PKP has emerged as a promising alternative to bilateral PKP and has been increasingly performed by doctors in recent years (5-12). However, whether unilateral PKP for OVCFs is more beneficial than bilateral PKP is still uncertain.

Meta-analysis of RCTs is considered be the best available source for evidence (36), Decision makers usually refer to meta-analyses to make recommendations on certain clinical topics. In light of this, an increasing number of meta-analyses have been reported to evaluate the differences between unilateral and bilateral PKP. However, discordant results have been found (33-38). These conflicting findings complicate decision

making depending on the best available evidence. Systematic reviews of discordant meta-analyses help to interpret and choose the currently discordant available evidence and are increasingly published in medical fields to facilitate decision-making (36,40-43). To the best of our knowledge, this is the first systematic review of discordant meta-analyses regarding unilateral PKP versus bilateral PKP for OVCFs.

Although the computer search of the meta-analyses was performed during a similar period, they did not comprise the same primary studies and did not reach the same conclusions for the treatment selection of OVCFs. The potential reasons for inconsistency amongst systematic reviews have been reported by Jadad et al (48), including the differences in clinical questions, study selection, eligibility criteria, abstracting data,





quality evaluation, the ability to pool studies, and data synthesis. Jadad et al (48) designed a decision algorithm to identify high-quality evidence from conflicting meta-analyses. This tool was reported to identify the best evidence among discordant meta-analyses (36,40-43).

On the basis of the Jadad decision algorithm, the meta-analysis reported by Feng et al (5) was chosen in the present study. Their study proved that unilateral PKP had a better degree of pain relief than bilateral PKP at post-operative short-term follow-ups. The radiological outcomes at post-operative short-term follow-ups suggested that bilateral PKP provided a better restoration of anterior vertebral height. Surgical time and cement dosage were lower when unilateral PKP was performed. No differences were found in complications between the groups, such as cement leakage and adjacent vertebral fractures. The findings of life quality suggested that unilateral PKP was better for bodily pain relief and provided general health benefits at post-operative short-term follow-ups. Therefore, we suggest that unilateral PKP may be superior to bilateral PKP for the treatment of OVCFs.

Although this study was strictly conducted following similar publications, several potential limitations may impair the power of the findings. First, this study only included articles in English. Some important articles in other languages were not included. The language bias is inevitable in this systematic review. Second, the primary RCTs had some defects in the methodologies including allocation concealment, inclusion of patients, and outcome assessment, which may result in overestimation of the advantages and/or disadvantages of both procedures.

## CONCLUSIONS

Based on this systematic review of discordant meta-analyses, the most current evidence suggests that unilateral PKP may be superior to bilateral PKP in the treatment of OVCFs. This study may aid doctors in evidence-based decisions regarding surgical selection of OVCFs treatment. However, high-quality studies should be conducted in the future because of the limitations of the studies included in this review.



REFERENCES

1. Johnell O, Kanis JA. An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporos Int* 2006; 17:1726-1733.
2. Yan L, Jiang R, He B, Liu T, Hao D. A comparison between unilateral transverse process-pedicle and bilateral puncture techniques in percutaneous kyphoplasty. *Spine (Phila Pa 1976)* 2014; 39:B19-B26.
3. Garfin SR, Buckley RA, Ledlie J, Balloon Kyphoplasty Outcomes Group. Balloon kyphoplasty for symptomatic vertebral body compression fractures results in rapid, significant, and sustained improvements in back pain, function, and quality of life for elderly patients. *Spine (Phila Pa 1976)* 2006; 31:2213-2220.
4. Rousing R, Hansen KL, Andersen MO, Jespersen SM, Thomsen K, Lauritsen JM. Twelve-months follow-up in forty-nine patients with acute/semiacute osteoporotic vertebral fractures treated conservatively or with percutaneous vertebroplasty: A clinical randomized study. *Spine (Phila Pa 1976)* 2010; 35:478-482.
5. Feng H, Huang P, Zhang X, Zheng G, Wang Y. Unilateral versus bilateral percutaneous kyphoplasty for osteoporotic vertebral compression fractures: A systematic review and meta-analysis of RCTs. *J Orthop Res* 2015; 33:1713-1723.
6. Sun H, Lu PP, Liu YJ, Yang X, Zhou PH, Shen XF, Sun SW, Yang H. Can unilateral kyphoplasty replace bilateral kyphoplasty in treatment of osteoporotic vertebral compression fractures? A systematic review and meta-analysis. *Pain Physician* 2016; 19:551-563.
7. Cheng X, Long HQ, Xu JH, Huang YL, Li FB. Comparison of unilateral versus bilateral percutaneous kyphoplasty for the treatment of patients with osteoporosis vertebral compression fracture (OVCF): A systematic review and meta-analysis. *Eur Spine J* 2016; 25:3439-3449.
8. Li LH, Sun TS, Liu Z, Zhang JZ, Zhang Y, Cai YH, Wang H. Comparison of unipedicular and bipedicular percutaneous kyphoplasty for treating osteoporotic vertebral compression fractures: A meta-analysis. *Chin Med J (Engl)* 2013; 126:3956-3961.
9. Huang Z, Wan S, Ning L, Han S. Is unilateral kyphoplasty as effective and safe as bilateral kyphoplasties for osteoporotic vertebral compression fractures? A meta-analysis. *Clin Orthop Relat Res* 2014; 472:2833-2842.
10. Yang LY, Wang XL, Zhou L, Fu Q. A systematic review and meta-analysis of randomized controlled trials of unilateral versus bilateral kyphoplasty for osteoporotic vertebral compression fractures. *Pain Physician* 2013; 16:277-290.
11. Lin J, Zhang L, Yang HL. Unilateral versus bilateral balloon kyphoplasty for osteoporotic vertebral compression fractures. *Pain Physician* 2013; 16:447-453.
12. Chen H, Tang P, Zhao Y, Gao Y, Wang Y. Unilateral versus bilateral balloon kyphoplasty in the treatment of osteoporotic vertebral compression fractures. *Orthopedics* 2014; 37:e828-e835.
13. Feng J ZQ, Xiao SX, Xia P. Clinical comparison of unilateral and bilateral percutaneous kyphoplasty for treating osteoporotic vertebral compression fracture. *Chinese J Trad Med Traum and Orthop* 2012; 20:26-29.
14. Li GZ WQ, Li S, Kang JP, Wang GJ. Clinical comparison study between unilateral and bilateral PKP for osteoporotic thoracolumbar compression fractures. *Orthopaedic Biomechanics Materials and Clinical Study* 2012; 9:17-21.
15. Chung HJ, Chung KJ, Yoon HS, Kwon IH. Comparative study of balloon kyphoplasty with unilateral versus bilateral approach in osteoporotic vertebral compression fractures. *Int Orthop* 2008; 32:817-820.
16. Chen C, Wei H, Zhang W, Gu Y, Tang G, Dong R, Xu Y, Chen L. Comparative study of kyphoplasty for chronic painful osteoporotic vertebral compression fractures via unipedicular versus bipedicular approach. *J Spinal Disord Tech* 2011; 24:E62-E65.
17. Huang S XQ, Xiang YC. Comparative study of unipedicular versus bipedicular percutaneous kyphoplasty for the treatment of osteoporotic vertebral compression fracture. *Orthopedic Journal of China* 2013; 21:115-118.
18. Feng J DW, Wang D. Comparison about the clinical effect of unilateral pedicle side kyphoplasty and bilateral transpedicular kyphoplasty for the patients with osteoporotic vertebral compression fractures. *China Medical Herald* 2013; 10:60-64.
19. YX H. Comparison of unilateral versus bilateral balloon kyphoplasty for osteoporotic vertebral compression fractures. *Chinese Journal of Clinical Rational Drug Use* 2012; 5:134-135.
20. Zhai HL ZF. Comparison of unipedicular and bipedicular balloon kyphoplasty for the treatment of osteoporotic vertebral compression fractures. *Inner Mongolia Medical J* 2013; 45:1501-1502.
21. Rebolledo BJ, Gladnick BP, Unnanuntana A, Nguyen JT, Kepler CK, Lane JM. Comparison of unipedicular and bipedicular balloon kyphoplasty for the treatment of osteoporotic vertebral compression fractures: A prospective randomised study. *Bone Joint J* 2013; 95-B:401-406.
22. Gu XH ZZ, Wu J, Lv J, Wu XY. Contrast study between unilateral and bilateral percutaneous balloon kyphoplasty for osteoporotic thoracolumbar compression fractures. *Chin J Surg Integr Tradit West Med (Chin)* 2009; 15:246-249.
23. He CJ YA. Effects of unipedicular versus bipedicular kyphoplasty for treatment of osteoporosis vertebral compression fractures. *Chin J Bone Joint Injury* 2014; 29:227-229.
24. Jiang Y SH, Wang D. The efficacy of unipedicular versus bipedicular percutaneous kyphoplasty for the treatment of osteoporotic vertebral compression fracture. *Chinese Journal of Bone and Joint Injury* 2010; 25:1011-1012.
25. Chen C, Chen L, Gu Y, Xu Y, Liu Y, Bai X, Zhu X, Yang H. Kyphoplasty for chronic painful osteoporotic vertebral compression fractures via unipedicular versus bipedicular approach: A comparative study in early stage. *Injury* 2010; 41:356-359.
26. XJ L. The randomized controlled study of unilateral versus bilateral percutaneous kyphoplasty for the treatment of elderly osteoporotic vertebral compression fractures. *J Cervicod Lumbod* 2014; 35:266-270.
27. Lin XL JW. Short and medium term observation of kyphoplasty for chronic osteoporotic vertebral compression fractures via unipedicular or bipedicular approach. *China Medical Herald* 2014; 11:34-40.
28. Chen L, Yang H, Tang T. Unilateral versus bilateral balloon kyphoplasty for multilevel osteoporotic vertebral compression fractures: A prospective study. *Spine (Phila Pa 1976)* 2011; 36:534-540.
29. Mao JG XM, Liu MQ. Unilateral versus bilateral balloon kyphoplasty for osteoporotic vertebral compression fractures. *Yiyao Qianyan* 2013; 23:20-21.
30. Zhang B DM, Tang Y. Unilateral versus bilateral kyphoplasty for osteopo-

- rotic vertebral compression fractures. *Advanced Materials Research* 2012; 393:1064-1068.
31. Li Q LM, Zheng CK. Unilateral versus bilateral percutaneous kyphoplasty for the treatment of osteoporotic compression fracture: A prospective study. *Medical Journal of Wuhan University* 2012; 33:567-570.
  32. Luo XL ZC, Kan WS, Li P. Unilateral versus bilateral percutaneous kyphoplasty for the treatment of osteoporotic compression fractures. *Chin J Tissue Eng Res* 2012; 16:567-570.
  33. Zhang H, Tang H, He Q, Wei Q, Tong D, Wang C, Wu D, Wang G, Zhang X, Ding W, Li D, Ding C, Liu K, Ji F. Surgical versus conservative intervention for acute achilles tendon rupture: A PRISMA-Compliant systematic review of overlapping meta-analyses. *Medicine (Baltimore)* 2015; 94:e1951.
  34. Ding F, Jia Z, Zhao Z, Xie L, Gao X, Ma D, Liu M. Total disc replacement versus fusion for lumbar degenerative disc disease: A systematic review of overlapping meta-analyses. *Eur Spine J* 2017; 26:806-815.
  35. Li Q, Wang C, Huo Y, Jia Z, Wang X. Minimally invasive versus open surgery for acute Achilles tendon rupture: A systematic review of overlapping meta-analyses. *J Orthop Surg Res* 2016; 11:65.
  36. Song GM, Bian W, Zeng XT, Zhou JG, Luo YQ, Tian X. Laparoscopic cholecystectomy for acute cholecystitis: Early or delayed?: Evidence from a systematic review of discordant meta-analyses. *Medicine (Baltimore)* 2016; 95:e3835.
  37. Wu Y, Lin L, Li H, Zhao Y, Liu L, Jia Z, Wang D, He Q, Ruan D. Is surgical intervention more effective than non-surgical treatment for acute Achilles tendon rupture? A systematic review of overlapping meta-analyses. *Int J Surg* 2016; 36:305-311.
  38. Zhao JG, Meng XH, Liu L, Zeng XT, Kan SL. Early functional rehabilitation versus traditional immobilization for surgical Achilles tendon repair after acute rupture: A systematic review of overlapping meta-analyses. *Sci Rep* 2017; 7:39871.
  39. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. *PLoS Med* 2009; 6:e1000100.
  40. Mascarenhas R, Saltzman BM, Sayegh ET, Verma NN, Cole BJ, Bush-Joseph C, Bach BR, Jr. Bioabsorbable versus metallic interference screws in anterior cruciate ligament reconstruction: A systematic review of overlapping meta-analyses. *Arthroscopy* 2015; 31:561-568.
  41. Erickson BJ, Mascarenhas R, Sayegh ET, Saltzman B, Verma NN, Bush-Joseph CA, Cole BJ, Bach BR Jr. Does operative treatment of first-time patellar dislocations lead to increased patellofemoral stability? A systematic review of overlapping meta-analyses. *Arthroscopy* 2015; 31:1207-1215.
  42. Zhao JG, Wang J, Wang C, Kan SL. Intra-medullary nail versus plate fixation for humeral shaft fractures: A systematic review of overlapping meta-analyses. *Medicine (Baltimore)* 2015; 94:e599.
  43. Zhao JG, Wang J, Long L. Surgical versus conservative treatments for displaced midshaft clavicular fractures: A systematic review of overlapping meta-analyses. *Medicine (Baltimore)* 2015; 94:e1057.
  44. Wright JG, Swiontkowski MF, Heckman JD. Introducing levels of evidence to the journal. *J Bone Joint Surg Am* 2003; 85-A:1-3.
  45. Shea BJ, Grimshaw JM, Wells GA, Boers M, Andersson N, Hamel C, Porter AC, Tugwell P, Moher D, Bouter LM. Development of AMSTAR: A measurement tool to assess the methodological quality of systematic reviews. *BMC Med Res Methodol* 2007; 7:10.
  46. Shea BJ, Hamel C, Wells GA, Bouter LM, Kristjansson E, Grimshaw J, Henry DA, Boers M. AMSTAR is a reliable and valid measurement tool to assess the methodological quality of systematic reviews. *J Clin Epidemiol* 2009; 62:1013-1020.
  47. Shea BJ, Bouter LM, Peterson J, Boers M, Andersson N, Ortiz Z, Ramsay T, Bai A, Shukla VK, Grimshaw JM. External validation of a measurement tool to assess systematic reviews (AMSTAR). *PLoS One* 2007; 2:e1350.
  48. Jadad AR, Cook DJ, Browman GP. A guide to interpreting discordant systematic reviews. *CMAJ* 1997; 156:1411-1416.