

Systematic Review

Evaluation of Sacroiliac Joint Interventions: A Systematic Appraisal of the Literature

Matthew P. Rupert, MD, MS¹, Marion Lee, MD², Laxmaiah Manchikanti, MD³,
Sukdeb Datta, MD⁴, and Steven P. Cohen, MD⁵

From: ¹Vanderbilt Interventional Pain Center Cool Springs, Cool Springs Surgery Center, Franklin, TN; ²The Pain Center at Affinity Health Group, Tifton, GA; ³Pain Management Center of Paducah, Paducah, KY; ⁴Vanderbilt University Medical Center, Nashville, TN; and ⁵Johns Hopkins School of Medicine, Baltimore, MD

Dr. Rupert is Assistant Professor of Interventional Pain Management, Vanderbilt Interventional Pain Center Cool Springs, Cool Springs Surgery Center, Franklin, TN. Dr. Lee is Director of The Pain Center at Affinity Health Group, Tifton, GA. Dr. Manchikanti is Medical Director of the Pain Management Center of Paducah, Paducah, KY, and Associate Clinical Professor of Anesthesiology and Perioperative Medicine, University of Louisville, Louisville, KY. Dr. Datta is Director, Vanderbilt University Interventional Pain Program, Associate Professor, Dept. of Anesthesiology, Vanderbilt University Medical Center, Nashville, TN. Dr. Cohen is Associate Professor, Department of Anesthesiology and Critical Care Medicine, Pain Management Division, Johns Hopkins School of Medicine, Baltimore, MD, and Walter Reed Army Medical Center, Washington, DC.

Address correspondence:
Matthew P. Rupert, MD, MS
Vanderbilt Interventional Pain Center
Cool Springs
Cool Springs Surgery Center
2009 Mallory Lane, Suite 250
Franklin, TN 37067
E-mail: mattrupert@comcast.net

Disclaimer: Dr. Datta receives research support from Sucampo Pharmaceuticals and an honorarium from Smith and Nephew.
Conflict of interest: None.

Manuscript received: 01/18/2008
Accepted for publication: 01/25/2009

Free full manuscript:
www.painphysicianjournal.com

Background: The sacroiliac joint has been implicated as a source of low back and lower extremity pain. There are no definite historical, physical, or radiological features that can definitively establish a diagnosis of sacroiliac joint pain. Based on the present knowledge, an accurate diagnosis is made only by controlled sacroiliac joint diagnostic blocks. The sacroiliac joint has been shown to be a source of pain in 10% to 27% of suspected patients with chronic low back pain utilizing controlled comparative local anesthetic blocks.

Study Design: A systematic review of diagnostic and therapeutic sacroiliac joint interventions.

Objective: To evaluate the accuracy of diagnostic sacroiliac joint interventions and the utility of therapeutic sacroiliac joint interventions.

Methods: The literature search was carried out by searching the databases of PubMed, EMBASE, and Cochrane reviews.

Methodologic quality assessment of included studies was performed using the Agency for Healthcare Research and Quality (AHRQ) methodologic quality criteria for diagnostic accuracy and observational studies, whereas randomized trials were evaluated utilizing the Cochrane review criteria. Only studies with scores of 50 or higher were included for assessment.

Level of evidence was based on the U.S. Preventive Services Task Force (USPSTF) criteria.

Outcome Measures: For diagnostic interventions, the outcome criteria included at least 50% pain relief coupled with a patient's ability to perform previously painful maneuvers with sustained relief using placebo-controlled or comparative local anesthetic blocks.

For therapeutic purposes, outcomes included significant pain relief and improvement in function and other parameters. Short-term relief for therapeutic interventions was defined as 6 months or less, whereas long-term effectiveness was defined as greater than 6 months.

Results: The indicated level of evidence is II-2 for the diagnosis of sacroiliac joint pain utilizing comparative, controlled local anesthetic blocks. The prevalence of sacroiliac joint pain is estimated to range between 10% and 38% using a double block paradigm in the study population. The false-positive rate of single, uncontrolled, sacroiliac joint injections is 20% to 54%.

The evidence for provocative testing to diagnose sacroiliac joint pain is Level II-3 or limited.

For radiofrequency neurotomy the indicated evidence is limited (Level II-3) for short- and long-term relief.

Limitations: The limitations of this systematic review include the paucity of literature evaluating the role of both diagnostic and therapeutic interventions and widespread methodological flaws.

Conclusions: The indicated evidence for the validity of diagnostic sacroiliac joint injections is Level II-2. The evidence for the accuracy of provocative maneuvers in the diagnosing of sacroiliac joint pain is limited (Level II-3). The evidence for radiofrequency neurotomy is also limited (Level II-3).

Key words: Chronic low back pain, sacroiliac joint pain, sacroiliitis, sacroiliac joint injection, sacroiliac joint dysfunction, thermal radiofrequency, pulsed radiofrequency

Pain Physician 2009; 12:399-418

Controlled studies have established intervertebral discs, facet joints, and sacroiliac joints as potential sources of low back and lower extremity pain (1-13). The sacroiliac joint is accepted as a potential source of low back and/or buttock pain with or without lower extremity pain. The sacroiliac joint has been implicated as the primary source of pain (1-8) in 10% to 27% (4,9,10) of patients with suspected sacroiliac joint pain utilizing controlled comparative local anesthetic blocks.

The sacroiliac joint is a diarthrodial joint, receiving innervation from the lumbosacral nerve roots (14-23). Neurophysiologic studies have demonstrated both nociceptive and proprioceptive afferent units in the sacroiliac joint (19,20,24). Referral patterns based on sacroiliac joint provocation and analgesic response to local anesthetics have been published in asymptomatic volunteers (25) and patients with pain (26,27).

There is no universally accepted gold standard for the diagnosis of low back pain stemming from the sacroiliac joint(s), intervertebral disc(s), or facet joint(s) (28). The recommended reference standards involve anesthetic or provocative injections (13). Multiple arguments have been made in favor and against the diagnostic accuracy of controlled local anesthetic blocks (1-8,12,28-35). However, controlled local anesthetic blocks continue to be the best available tool to identify either the intervertebral discs, facet, or sacroiliac joints as the source of low back pain (1-3,13,28). Yet, these reference standards are invasive, expensive, and often difficult to interpret, and therefore may not be suitable for routine clinical use.

In a systematic review evaluating a battery of tests to identify the disc, sacroiliac joint, or facet joint as the source of low back pain, Hancock et al (28) suggested that a combination of sacroiliac joint pain provocative maneuvers appears to be useful in pinpointing the sacroiliac joint as the principal source of symptoms in patients with pain below the 5th lumbar vertebra. They also concluded that although a positive bone scan has high specificity, it is associated with a very low sensitivity, which means that the majority of patients with the sacroiliac joint pain will not be accurately identified.

A systematic review by Szadek et al (36), evaluated the diagnostic validity of the International Association for the Study of Pain (IASP) criteria for sacroiliac joint pain. The meta-analysis showed that the thigh thrust test, the compression test, and 3 or more positive stressing tests contain sufficient discriminative

power for diagnosing sacroiliac joint pain. They concluded that in view of the lack of a gold standard for sacroiliac joint pain, the diagnostic validity of tests for sacroiliac joint pain should be regarded with caution.

Song et al (37) performed a systematic literature review evaluating the diagnostic value of scintigraphy in assessing sacroiliitis and ankylosing spondylitis. They concluded that scintigraphy is at best of limited value in establishing a diagnosis of ankylosing spondylitis. Two systematic reviews evaluated the role of diagnostic intraarticular injections (1,2) in establishing the sacroiliac joint(s) as the primary pain generator. Both reviews concluded the specificity and validity to be moderate. In a best-evidence review of diagnostic procedures for neck and low back pain that focused on previously published systematic reviews (1,2,28), Rubinstein and van Tulder (38) also concluded that there was moderate evidence for diagnostic sacroiliac joint blocks.

Sacroiliac joint pain may be managed by intraarticular injections or neurolysis of the nerve supply. However, 2 previous systematic reviews (1,2) found the evidence supporting therapeutic sacroiliac joint interventions to be limited. European guidelines for the management of chronic non-specific low back pain (39) evaluating the literature through 2002 also concluded that there is limited evidence supporting sacroiliac joint injections with corticosteroids. But despite the absence of any clear consensus in favor of sacroiliac joint interventions, their use has continued to grow in recent years (40-42).

The purpose of this review is to systematically assess the literature on diagnostic and therapeutic sacroiliac joint interventions.

METHODS

The literature search included the databases PubMed, EMBASE, and Cochrane reviews; systematic and narrative reviews; and the NIH clinical trials registry. The search included articles published between 1966 and 2008. A manual review of the reference section of selected articles was then performed to identify relevant studies missed in the electronic search. Only English language articles were reviewed.

The search was conducted utilizing the following terms: sacroiliac joint, sacroiliac joint pain, sacroiliac joint injections, radiofrequency neurotomy of sacroiliac joint, neurolytic blocks of sacroiliac joint.

Diagnostic Sacroiliac Joint Interventions

Inclusion Criteria

Diagnostic sacroiliac joint interventions in patients with chronic low back and/or lower extremity pain for greater than 3 months in duration were analyzed for review. Only those studies utilizing fluoroscopically guided controlled diagnostic blocks (i.e., placebo-controlled or comparative local anesthetic) were included. The criterion standard for diagnosis of sacroiliac joint pain was 50% or greater pain relief for the duration of action of the local anesthetic, coupled with increased ability to perform previously painful movements.

Excluded from analysis were studies done on animals, cadavers, ultrasound guided injections, case reports, book chapters, non-evidence-based guidelines, letters to the editor, and expert opinion papers.

Method of Review

All abstracts obtained from computerized database searches were screened for inclusion. Two physician reviewers evaluated articles meeting inclusion criteria for methodologic quality using the modified Agency for Healthcare Research and Quality (AHRQ) criteria for diagnostic studies (43).

The quality of individual articles was evaluated using the above criteria with application of consensus-based weighted scores developed by the guidelines committee of the American Society of Interventional Pain Physicians (ASIPP) (33). These guidelines have been utilized in other systematic reviews (44,45) and have recently been refined and modified (46-50). Only studies scoring at least 50 out of 100 were included

for analysis.

Each study was evaluated by 2 physicians for the stated criteria with any disagreements resolved by a third physician. If there was a conflict of interest with the reviewed manuscripts with authorship or any other type of conflict, the involved authors did not review the manuscripts for quality assessment, clinical relevance, evidence synthesis, or grading of evidence.

Analysis of Evidence

Qualitative analysis was conducted using 5 levels of evidence, ranging from Level I to III with 3 subcategories in Level II, as illustrated in Table 1 (51).

Therapeutic Sacroiliac Joint Interventions

Inclusion Criteria

Studies should have documented the existence of sacroiliac joint pain using controlled sacroiliac joint blocks. Two types of SI joint interventions were included in this review: intraarticular sacroiliac joint injections and radiofrequency neurotomy of the nerve supply to the sacroiliac joint. All studies must have documented outcome evaluations extending at least 6 months, with appropriate statistical analysis.

Studies done without appropriate diagnostic methods (i.e., minimizing false-positive responses, non-systematic reviews, book chapters, and case reports) were excluded.

Outcome Parameters

The primary outcome measure was pain relief at various time points documented over a period lasting

Table 1. *Modified quality of evidence developed by USPSTF.*

I:	Evidence obtained from multiple properly conducted diagnostic accuracy studies.
II-1:	Evidence obtained from at least one properly conducted diagnostic accuracy study of adequate size.
II-2:	Evidence obtained from at least one properly designed small diagnostic accuracy study.
II-3:	Evidence obtained from diagnostic studies of uncertainty.
III:	Opinions of respected authorities, based on clinical experience descriptive studies and case reports or reports of expert committees.

Adapted from the U.S. Preventive Services Task Force (USPSTF) (51).

at least 6 months. Secondary outcome measures were functional improvement, psychological improvement, return-to-work, opioid use, and complications. Short-term relief was defined as relief lasting 6 months or less and long-term relief as benefits extending beyond 6 months.

Methodologic Quality Assessment

The quality of each individual article used in this analysis was assessed by modified Cochrane review criteria with weighted scores (52) for randomized trials and AHRQ quality criteria for assessment for observational studies (43) with consensus-based weighted scoring developed by the guidelines committee of ASIPP (33) and used in other systematic reviews (44,47-50,53-61). Only studies scoring at least 50 out of 100 were included for analysis.

Each study was evaluated by 2 physicians for the stated criteria with any disagreements resolved by a third physician. If there was a conflict of interest with the reviewed manuscripts with authorship or any other type of conflict, the involved authors did not re-

view the manuscripts for quality assessment, clinical relevance, evidence synthesis, or grading of evidence.

Analysis of Evidence

Qualitative analysis was conducted using 5 levels of evidence, ranging from Level I to III with 3 subcategories in Level II, as illustrated in Table 1 (51).

Recommendations

Grading recommendations were based on Guyatt et al's criteria with 6 Levels, 1A – 1C strong and 2A – 2C weak as illustrated in Table 2 (62).

RESULTS

Diagnostic Studies

Literature Search


Our comprehensive search yielded 2,260 articles for review on sacroiliac joint pain (Fig. 1). However, only 13 studies were considered for inclusion.

The following studies were excluded for failure

Table 2. Grading recommendations.

Grade of Recommendation/ Description	Benefit vs Risk and Burdens	Methodological Quality of Supporting Evidence	Implications
1A/strong recommendation, high-quality evidence	Benefits clearly outweigh risk and burdens, or vice versa	RCTs without important limitations or overwhelming evidence from observational studies	Strong recommendation, can apply to most patients in most circumstances without reservation
1B/strong recommendation, moderate quality evidence	Benefits clearly outweigh risk and burdens, or vice versa	RCTs with important limitations (inconsistent results, methodological flaws, indirect, or imprecise) or exceptionally strong evidence from observational studies	Strong recommendation, can apply to most patients in most circumstances without reservation
1C/strong recommendation, low-quality or very low-quality evidence	Benefits clearly outweigh risk and burdens, or vice versa	Observational studies or case series	Strong recommendation but may change when higher quality evidence becomes available
2A/weak recommendation, high-quality evidence	Benefits closely balanced with risks and burden	RCTs without important limitations or overwhelming evidence from observational studies	Weak recommendation, best action may differ depending on circumstances or patients' or societal values
2B/weak recommendation, moderate-quality evidence	Benefits closely balanced with risks and burden	RCTs with important limitations (inconsistent results, methodological flaws, indirect, or imprecise) or exceptionally strong evidence from observational studies	Weak recommendation, best action may differ depending on circumstances or patients' or societal values
2C/weak recommendation, low-quality or very low-quality evidence	Uncertainty in the estimates of benefits, risks, and burden; benefits, risk, and burden may be closely balanced	Observational studies or case series	Very weak recommendations; other alternatives may be equally reasonable

Adapted from Guyatt G et al. Grading strength of recommendations and quality of evidence in clinical guidelines. Report from an American College of Chest Physicians task force. *Chest* 2006; 129:174-181 (62).



to exclude patients with comparative blocks or unresolvable technical flaws (Table 3).

Fortin et al (25) used pain provocation during a single anesthetic block. Schwarzer et al (11) used $\geq 75\%$ reduction in pain following a single injection of local anesthetic in patients with pain experienced below L5/S1. Dreyfuss et al (63) used a single injection of local anesthetic and corticosteroid, noted pain provocation, and required more than 90% diminution in the mean pain score. Slipman et al (27) used $\geq 80\%$ reduction in general pain on a vascular autonomic signal (VAS) following a single anesthetic injection in low back pain patients. Laslett et al (64) and Young et al (65) used greater than 80% relief as their criterion standard following single blocks. Maigne et al (66) and Slipman et al (67) both utilized single diagnostic blocks with 75% and 80% pain relief, respectively, in evaluating the sensitivity and specificity of radionuclide bone scanning. Broadhurst and Bond (68) utilized a placebo-controlled injection with $\geq 70\%$ relief and the reduction of pain provocation. However, this study had multiple issues relating to the technical aspects of the injection and volume of injectate (69). Pang et al (70) also utilized a single block. Maigne and Planchon (12) evaluated sacroiliac joint pain after lumbar fusion with a single block with a criterion standard of 75% pain relief on a VAS with 35% positive rate of blocks.

Multiple authors used dual blocks with 2 local anesthetics of different duration of action (4,9,10,71-73). These studies met all inclusion criteria for diagnostic accuracy evaluation and thus underwent methodological quality assessment.

Diagnostic Accuracy Studies

Five studies were incorporated for methodologic quality assessment of diagnostic accuracy of diagnostic sacroiliac joint injections (4,9,10,71-73).

Methodologic Quality Assessment

The methodologic quality assessment of the 5 studies meeting inclusion criteria are illustrated in Table 4. Two publications by van der Wurff et al (72,73) were from a single study; consequently, 5 studies were included in the methodologic quality assessment. The series of blocks in these studies were performed under fluoroscopic guidance using lidocaine and bupivacaine, with the response and criterion being at least 50% pain relief.

Study Characteristics

Table 5 shows the characteristics of the diagnostic studies evaluating the accuracy and prevalence.

Maigne et al (9) studied 54 patients who had chronic (> 50 days) unilateral low back pain (VAS > 4) with or without radiation to the posterior thigh with associated pain and tenderness over the posterior sacroiliac joint. Dual injections were performed, first with a screening lidocaine injection (2 mL), then with bupivacaine. Nineteen of 54 patients had $\geq 75\%$ relief from the screening block and 10 of 19 participants had $\geq 75\%$ improvement lasting longer than 2 hours from the confirmatory bupivacaine block. Among the 54 subjects, 10 or 18.5% were considered to have sacroiliac joint pain. The false-positive rate using the total number of screening blocks was determined to be

Table 3. Illustration of studies excluded.

Study	Reason for Exclusion
Fortin et al (25)	Used pain provocation during a single anesthetic block.
Schwarzer et al (11)	Used single injection of local anesthetic.
Dreyfuss et al (63)	Used a single injection of local anesthetic and corticosteroid.
Slipman et al (27)	Single anesthetic injection in low back pain patients was used.
Laslett et al (64)	Single blocks were used.
Young et al (65)	Single blocks were used.
Maigne et al (66)	Utilized single diagnostic blocks.
Slipman et al (67)	Utilized single diagnostic blocks.
Broadhurst and Bond (68)	This study had multiple issues relating to the technical aspects of the injection and volume of injectate (71).
Pang et al (70)	Utilized a single block.
Maigne and Planchon (12)	Evaluated with a single block.

Table 4. Methodological quality assessment and scoring of sacroiliac joint studies.

STUDY	Study Population (15)	Adequate Description of Test (10)	Appropriate Reference Standard (30)		Blinded Comparison of Test (30)		Avoidance of Verification Bias (15)	TOTAL (100)
			Appropriate reference standard (gold standard) used for comparison (15)	Reference standard reproducible (15)	Evaluation of test without knowledge of disease status, if possible (15)	Independent, blind interpretation of test and reference (15)		
Manchikanti et al (4)	15	10	5	5	5	10	15	65
Maigne et al (9)	15	10	5	5	5	10	15	65
Irwin et al (10)	15	10	5	5	5	10	15	65
Laslett et al (71)	15	10	5	5	5	10	15	65
van der Wurff et al (72)	15	10	5	5	5	10	15	65

() weighted item score

Methodological criteria and scoring adapted from West S et al. Systems to Rate the Strength of Scientific Evidence, Evidence Report, Technology Assessment No. 47. AHRQ Publication No. E016 (43).

20% (12). Since only patients with a high likelihood of sacroiliac joint pain were tested, the calculated prevalence rate of 18.5% may not be reliable. The authors also excluded 3 patients who remained pain-free after receiving the screening block. In addition, 3 others had a temporary sciatic palsy after the first block resulting from leakage of the anesthetic. In 7 cases, penetration of the sacroiliac joint was technically impossible because of degenerative changes. Consequently, among the 67 enrolled patients, only 54 completed the study because the injected contrast was insufficient to fill the entire joint cavity, no arthrographic interpretation was conducted.

Manchikanti et al (4) evaluated 120 patients who presented to a pain clinic with low back pain of ≥ 6 months. Twenty patients with suspected sacroiliac joint involvement (i.e., negative facet blocks, spontaneous pain in the sacral region, sacroiliac joint tenderness, and positive provocative maneuvers) then underwent screening sacroiliac joint injections with 2% lidocaine followed in 3 to 4 weeks by confirmatory bupivacaine blocks. Six patients had $\geq 80\%$ short-acting pain relief following the screening block. Among these subjects, 2 patients experienced concordant pain relief after the confirmatory bupivacaine block. When calculated for all patients with low back pain, the overall prevalence

of SI joint pain was 2%. When patients with radicular symptomatology and facetogenic pain were ruled out, the point prevalence rate increased to 10% with a false-positive rate of 22%. Despite the otherwise strong methodology, the small sample size limits the applicability and conclusions that can be drawn.

In a retrospective review involving 158 patients, Irwin et al (10) evaluated the prevalence of sacroiliac joint pain in various subgroups using dual comparative local anesthetic blocks. Fluoroscopically guided intraarticular injections were performed first with 2 mL of 2% lidocaine, followed by 2 mL of 0.25% bupivacaine with corticosteroid for the confirmatory injection. A positive response was defined as greater than 70% pain relief lasting at least 3 and 4 hours after the first and second injections, respectively. Overall, the percentage of patients diagnosed with sacroiliac joint pain using the dual comparative local anesthetic injections was 26.7% (42/158). The authors found that patients diagnosed with sacroiliac joint pain tend to be older than those with other pain sources, whereas, gender, age, and smoking status were not correlated with the response to diagnostic blocks. Although this study was the largest and one of the most externally valid among the studies analyzed, its retrospective nature raises multiple issues related to validity. These issues include

Table 5. Characteristics of reported diagnostic studies evaluating prevalence.

Study	Participants	Objective(s)	Interventions(s)	Result(s)
Maigne et al (9)	54 patients aged 18-75 with chronic unilateral LBP with or without radiation to the posterior thigh for > 50 days (median 4.2 months). Patients had failed epidural or lumbar facet injections.	To determine the prevalence of sacroiliac joint pain in a selected population of patients with low back pain and assess certain pain provocation tests.	Successful blockade of the sacroiliac joint in 54 patients. A screening block was done with 2% lidocaine and a confirmatory block was performed with bupivacaine 0.5%. Greater than 75% relief was considered a positive block.	Prevalence = 18.5% False-positive rate = 20%
Manchikanti et al (4)	120 patients (age 18-90) presenting to the clinic with > 6 months of low back pain and no structural basis for the pain by radiographic imaging. 20 patients were evaluated for SI joint pain.	To determine the frequency of various structures responsible for low back pain.	All patients had facet blocks. Non-responders who fit criteria had double injection sacroiliac joint blocks. The screening block was done with 2% lidocaine and the confirmatory block was performed using 0.5% bupivacaine.	The incidence of sacroiliac joint pain was 2% of the overall sample and 10% of those suspected to have sacroiliac joint pain. The false-positive rate was 22%.
Irwin et al (10)	158 patients underwent sacroiliac joint injections with average symptoms duration of 34 months. Patients failed conservative modalities prior to injection therapy.	To evaluate prevalence and correlation between age, gender, and body mass index by dual comparative local anesthetic blocks.	The fluoroscopically guided contrast-enhanced sacroiliac joint injections were performed initially with 2 mL of 2% lidocaine for the first injection, followed by 2 mL of 0.25% bupivacaine, a local anesthetic, for the confirmatory injection. A patient was required to have at least 70% reduction of familiar painful symptoms after the initial injection for 3 or 4 hours for positive response.	26.6% were found to have sacroiliac joint pain by dual injections. Estimated false-positive rate = 53.8%
Laslett et al (71)	48 patients received initial sacroiliac joint diagnostic injection, derived from 62 patients with buttock pain with or without lumbar or lower extremity symptoms.	To assess the diagnostic accuracy of clinical examination in identifying symptomatic and asymptomatic sacroiliac joints using double-diagnostic injections as reference standard.	16 patients had positive response to sacroiliac joint injections and 5 of them did not receive a confirmatory diagnostic injection because they derived such symptomatic relief from the initial procedure that a confirmatory injection could not be justified. 11 patients received a confirmatory injection and all of them tested positive. Overall 32 patients had negative sacroiliac joint injections and did not require a confirmatory injection.	25.6% were found to have sacroiliac joint pain by dual injections. Estimated false-positive rate = none
van der Wurff et al (72)	Total number of 140 patients with chronic low back pain visiting the pain clinic in the Netherlands, 60 patients entered the study.	To compare the diagnostic accuracy of multi-test regimen of 5 sacroiliac joint pain provocation test with fluoroscopically controlled double sacroiliac joint blocks using a short- and long-acting local anesthetic.	The fluoroscopically guided contrast enhanced sacroiliac joint injections were performed initially with 2 mL of 2% lidocaine and next time with 0.25% bupivacaine. A reduction in the patient's characteristic pain of 50% or more on the VAS remaining for at least one hour for lidocaine or 4 hours for bupivacaine was considered as positive. When a patient showed a VAS reduction after both intraarticular sacroiliac joint blocks, this was considered a positive response. Any other outcome was considered a negative response.	Prevalence = 38% False-positive rate = 21%

non-uniform screening procedures, and lack of standardization regarding injection technique. The higher false-positive rate reported in this study (53.8%) can be attributed to using the number of positive screening blocks as the denominator, rather than the initial number of injections

van der Wurff et al (72) evaluated 60 patients meeting the inclusion criteria of pain below L5 overlying the posterior aspect of sacroiliac joint and pain intensity exceeding 50 on a 0 – 100 VAS scale. Excluded patients included those with ankylosing spondylitis, leg-length discrepancies greater than 2 cm, positive Waddell's signs, clinical osteoarthritis, and radicular pain with focal neurological signs or symptoms. Each patient underwent 2 diagnostic sacroiliac joint injections on separate occasions with short and long-acting local anesthetics. All blocks were performed by an anesthesiologist unaware of the results of provocation testing by an independent investigator who examined the patient prior to the procedure. A positive response was pre-designated as $\geq 50\%$ pain relief lasting for at least one hour after the intraarticular lidocaine injection and at least 4 hours after the bupivacaine block. Any other outcome was considered to be a negative response. Twenty-seven patients (45%) responded positively to both blocks. Among those patients with 3 or more positive provocative tests, 29 achieved a positive response following the initial block, with 23 obtaining prolonged pain relief after the bupivacaine injection (true-positives). One patient did not respond to either block.

Laslett et al (71) evaluated 48 patients who underwent screening diagnostic sacroiliac joint injections from amongst a sample of 62 with buttock pain with or without lumbar or lower extremity symptoms. Sixteen patients had a positive response, predetermined to be $\geq 80\%$ relief. Five responders obtained

long-term relief, leaving 11 patients who underwent confirmatory blocks with bupivacaine. All confirmatory blocks were positive. Ten of the 11 sacroiliac joint patients met clinical examination criteria for having sacroiliac joint pain, defined as 3 (of 5) positive provocation tests in absence of peripheralization or centralization. Potential sources of bias were that these patients were specifically selected for injection therapy and exposed to multiple procedures. Excluding the 5 patients with a prolonged response to the lidocaine screening blocks, the prevalence rate in this pre-selected was 26% (Table 6). Potential confounding factors include the lack of false-positive responses and that some of these patients were included in other studies and publications (64,65,71).

Level of Evidence

The indicated level of evidence is Level II-2, as illustrated in Table 1.

Provocation Testing

There is no universally accepted gold standard for the diagnosis of low back pain originating from the sacroiliac joint. The recommended reference standard involves either anesthetic or provocative injections; however, doubts have been cast on the validity of a sacroiliac joint block as a diagnostic gold standard. A review by Berthelot et al (74) concluded clinical signs and maneuvers to be unreliable for diagnosing pain originating within the sacroiliac joint, being fraught with both low sensitivity and specificity. But this review also concluded that sacroiliac joint blocks were similarly unreliable, since pain patterns formerly attributed to the sacroiliac joint can be related to extraarticular structures, most notably the numerous ligaments surrounding the joint.

Table 6. *Data of prevalence of sacroiliac joint pain by controlled diagnostic blocks.*

Study	Methodologic Criteria	# of Subjects	Prevalence Estimates	False-Positive Rate
Manchikanti et al (4)	65	20	10% (95% CI, 0% - 23%)	22% (95% CI, 3% - 42%)
Maigne et al (9)	65	54	18.5% (95% CI, 8% - 29%)	20% (95% CI, 8% - 33%)
Irwin et al (10)	65	158	26.6% (95% CI, 20% - 34%)	53.8% (95% CI, 43% - 64%)
Laslett et al (71)	65	43/48	25.6% (95% CI, 12% - 39%)	0%
van der Wurff et al (72)	65	60	38% (95% CI, 26% - 51%)	21% (95% CI, 7% - 35%)

CI = confidence interval

In contrast, Szadek et al (36) and Hancock et al (28) in 2 separate systematic reviews showed a positive correlation between provocation testing and diagnostic blocks. Hancock et al (28) included 6 studies in their analysis (63,64,66,67,68,71,72) while Szadek et al included 15 (36). In a systematic appraisal of the literature assessing the accuracy of multiple tests for back pain utilizing QUADAS criteria, Simpson and Gemmill (75) identified 5 studies that focused on sacroiliac joint pain (63,64,68,71,76). They found no single test to be consistently valid.

Among the studies analyzed in these reviews, 3 employed dual blocks (9,71,72), and hence were considered.

Maigne et al (9) determined the prevalence of sacroiliac joint pain in a selected population of patients suffering from low back pain and assessed the validity of various pain provocation tests. The patients underwent 7 sacroiliac pain provocation tests, which included the distraction test, compression test, sacral pressure test, Gaenslen's test, Patrick's test, resisted external rotation of the hip, and direct pressure on the pubic symphysis, before and after a screening block. However, no statistically significant association was found between the response to the blocks and any single clinical parameter. They concluded that no pain provocation test was a useful predictor of sacroiliac joint pain.

Laslett et al (71) concluded that the diagnostic accuracy of the clinical examination in conjunction with reasoning processes was superior to sacroiliac joint pain provocation maneuvers as stand alone tests. Excluding patients whose pain centralized or peripheralized increased the positive likelihood ratio for identifying a symptomatic sacroiliac joint(s) in patients with 3 or more provocative tests.

van der Wurff et al (72) evaluated the diagnostic accuracy of a multi-test regimen of 5 sacroiliac joint pain provocation tests by comparing it to the results of fluoroscopically guided double local anesthetic blocks. The five provocation tests included the distraction test, compression test, thigh thrust, Patrick's sign, and Gaenslen's test. Among the 60 patients studied, 45% obtained a positive response to both blocks. Whereas none of the provocation tools were specific as a stand-alone test, a combination of 3 or more positive tests was deemed to be a reliable indicator. Seven patients were categorized as having false-positive blocks. Four of the 23 double-block patients obtained complete relief of their pain after injection such that

19 patients could be considered "true-positives." The false-positive rate for this study was estimated to be 21% with a prevalence rate of 38%. The authors' conclusion that a correlation exists between the finding of 3 or more positive pain provocation tests and an analgesic response to double intraarticular sacroiliac joint blocks corroborates the results of Laslett et al (71). This study is limited by a pain relief criterion standard response of $\geq 50\%$ rather than $\geq 80\%$ pain relief after injection.

Level of Evidence

The indicated evidence for the accuracy of provocative maneuvers in diagnosis of sacroiliac joint injections pain is limited or Level II-3.

Accuracy of Imaging and Clinical History

The value of medical history, physical examination, and radiological imaging has been questioned in the diagnosis of sacroiliac joint pain. In a prospective study evaluating the diagnostic utility of a battery of accepted sacroiliac joint tests recommended by a multidisciplinary expert panel, Dreyfuss et al (63) attempted to discern whether a single test or an ensemble of investigative maneuvers were sufficiently useful in identifying a painful sacroiliac joint. They concluded that sacroiliac joint pain is resistant to identification by history and physical examination data. Although distinct pain patterns have been previously described by an assortment of investigators (11,25,27), similar referral maps may be produced by other structures. Schwarzer et al (11) found radiation to the groin to be the only descriptive quality reliably associated with pain relief after a single sacroiliac joint block.

Noting that history and physical examination findings associated with somatically referred pain can be unreliable and non-specific, Jung et al (77) evaluated the accuracy of "pain distribution pattern templates" in predicting pain relief following confirmatory intraarticular and deep interosseous ligament blocks. They concluded that 46% of patients with sacroiliac joint arthropathy could be diagnosed based on pain distribution patterns.

Neither has radiological imaging been found to be an accurate indicator of a painful sacroiliac joint. Puhakka et al (78,79) performed 2 evaluations, one with magnetic resonance imaging (MRI) of the normal sacroiliac joint with correlation to histology and the second one with MRI abnormalities of the sacroiliac

joints in early spondyloarthropathy with a one-year follow-up study. They concluded that coronal MRI does not allow assessment of normal anatomy. But when there are variants or abnormalities of the ventral and dorsal margins of the cartilaginous sacroiliac joint and in early spondyloarthropathy, MRI can detect significant inflammatory and destructive changes of the sacroiliac joints over a one-year follow-up, in spite of minimal changes in the clinical parameters. However, the MRI changes and inflammatory activity are not detectable by CT and x-ray examinations. Thus, MRI may be a sensitive method, without risks, for early diagnosis and for following disease progression in spondyloarthropathy. Radiologic studies can assist in determining anatomical integrity (80). A retrospective study (81) showed that CT scans were negative in 42% of symptomatic sacroiliac joints.

Song et al (37) evaluated the diagnostic value of scintigraphy in detecting sacroiliitis in ankylosing spondylitis and those with probable sacroiliitis without x-ray changes. Following an extensive literature search, they concluded that scintigraphy of the sacroiliac joints is at most of limited diagnostic value.

Therapeutic Sacroiliac Joint Interventions

The literature search yielded 33 relevant evaluations for intraarticular injections and radiofrequency neurotomy (Fig. 2).

Among the many studies considered for inclusion (82-109), there were 5 randomized trials (83,84,86,91,107), with the remainder consisting of retrospective reviews and observational reports (82,85,87-91,93-106,108,109).

Intraarticular Sacroiliac Joint Injections

Based on the search, 17 publications were selected (82-98). Of these, 4 studies were randomized trials (83,84,86,91) and 14 were observational reports (82,85,87-90,92-99).

Among the 4 randomized trials meeting criteria for initial evaluation (83,84,86,91), all were excluded due to lack of a valid diagnosis prior to therapeutic interventions. These studies also had other methodological issues such as lack of long-term follow-up (83), using children as the target population (86), and evaluating periarticular injections done for spondyloarthropathy (91).

Among the 14 observational reports (82,85,87-90,92-99), none met inclusion criteria. The basis for exclusion included lack of controlled diagnostic blocks to

establish diagnosis of sacroiliac joint pain, evaluating only patients with spondyloarthropathy (86,87,89,90), and not following patients for 6 months. Thus, no methodologic quality assessment was performed for the category of intraarticular injections.

Radiofrequency Neurotomy

The literature search yielded 9 relevant reports available for review (100-108). Of these, one was randomized (107) and the rest were observational.

Methodologic Quality Assessment


The study by Cohen et al (107) was double blind placebo-controlled; however, it utilized a single diagnostic block as the means for diagnosis. Another study (103) had inconsistent descriptions utilizing interligamentous rather than intraarticular injections. Ferrante et al (100), Gevarguez et al (101), and Buijs et al (104) all utilized single diagnostic blocks. Kapural et al (108) had only short-term follow-up of 3 to 4 months, even though they utilized dual blocks.

Burnham and Yasui (105), Cohen and Abdi (102), and Vallejo et al (106) utilized dual blocks.

Methodologic quality assessment is illustrated in Table 7.

Characteristics of Included Studies

Vallejo et al (106) tested the hypothesis that pulsed radiofrequency of the posterior rami from L4 to S3 would provide therapeutic benefit to patients with intractable sacroiliac joint dysfunction. One hundred and twenty-six patients with suspected sacroiliac joint pain were examined for this study. Dual diagnostic blocks with local anesthetic and corticosteroid using $\geq 75\%$ relief as the success criterion were done to minimize false-positive results and confirm the pain generator. This resulted in 52 patients with confirmed disease. Thirty of these patients obtained $\geq 50\%$ relief lasting longer than 12 weeks. The remaining 22 subjects were offered the treatment. The follow-up period was 6 months and outcome measures included VAS scoring and a quality of life assessment tool. Sixteen of the 22 were found to have good ($\geq 50\%$) to excellent ($\geq 80\%$) results; however, in only 7 patients did this improvement exceed 17 weeks. There was no annotation about how many patients obtained 6 or greater months of relief. This study is limited by its observational nature, and the small number of patients. In addition, only 7 of 22 patients experienced between 17 and 32 weeks worth of relief, which is similar to



Sacroiliac Joint Injections

Table 7. *Methodologic quality assessment criteria for radiofrequency neurotomy of sacroiliac joint.*

	Weighted Score (points)	Vallejo et al (106)	Cohen & Abdi (102)	Burnham & Yasui (105)
1. Study Question	2	2	2	2
• Clearly focused and appropriate question		2	2	2
2. Study Population	8	5	5	5
• Description of study population	5	5	5	5
• Sample size justification	3	0	–	–
3. Comparability of Subjects	22	5	5	5
• Specific inclusion/exclusion criteria for all groups	5	5	5	5
• Criteria applied equally to all groups	3	–	–	–
• Comparability of groups at baseline with regard to disease status and prognostic factors	3	–	–	–
• Study groups comparable to non-participants with regard to confounding factors	3	–	–	–
• Use of concurrent controls	5	–	–	–
• Comparability of follow-up among groups at each assessment	3	–	–	–
4. Exposure or Intervention	11	8	8	8
• Clear definition of exposure	5	5	5	5
• Measurement method standard, valid and reliable	3	3	3	3
• Exposure measured equally in all study groups	3	–	–	–
5. Outcome measures	20	13	14	15
• Primary/secondary outcomes clearly defined	5	5	5	5
• Outcomes assessed blind to exposure or intervention	5	–	–	–
• Method of outcome assessment standard, valid and reliable	5	5	5	5
• Length of follow-up adequate for question	5	3	4	5
6. Statistical Analysis	19	8	8	8
• Statistical tests appropriate	5	5	5	5
• Multiple comparisons taken into consideration	3	3	3	3
• Modeling and multivariate techniques appropriate	2	–	–	–
• Power calculation provided	2	–	–	–
• Assessment of confounding	5	–	–	–
• Dose-response assessment if appropriate	2	–	–	–
7. Results	8	5	5	6
• Measure of effect for outcomes and appropriate measure of precision	5	3	3	3
• Adequacy of follow-up for each study group	3	2	2	3
8. Discussion	5	5	5	5
• Conclusions supported by results with possible biases and limitations taken into consideration		5	5	5
9. Funding or Sponsorship	5	5	5	5
• Type and sources of support for study		5	5	5
TOTAL SCORE	100	56	57	59

Adapted and modified from West S et al. Systems to Rate the Strength of Scientific Evidence, Evidence Report, Technology Assessment No. 47. AHRQ Publication No. 02-E016 (43).

the duration of benefit obtained from local anesthetic blocks with or without steroids (110-116).

Burnham and Yasui (105) published the results of a pilot study evaluating bipolar radiofrequency neurotomy. They evaluated 9 subjects with sacroiliac joint pain confirmed by local anesthetic joint and lateral branch nerve blocks. These subjects were treated with a series of radiofrequency strip lesions performed adjacent to the lateral dorsal foraminal aperture plus conventional monopolar lesioning at the L5 dorsal ramus. Follow-up visits were conducted at one, 3, 6, 9, and 12 months after the procedure. Significant reductions in back and leg pain frequency and severity, and analgesic intake were demonstrated at all points. Complications were minimal. Overall, 8 of the 9 subjects were satisfied with the procedure. The median improvement in pain intensity was 4.1 on a 0 – 10 numeric rating scale and the reduction in disability was 17.8 on the Oswestry Disability Index (ODI). Overall satisfaction was 67% at 12 month follow-up. Limitations include the small number of patients (n = 9) recruited from one practice.

Cohen and Abdi (102) performed radiofrequency lesioning on 9 patients who experienced greater than 80% pain relief following intraarticular joint injection(s) and greater than 50% relief following L4-5 primary dorsal rami and S1-3 lateral branches blocks. Eight of 9 patients (89%) obtained 50% or greater pain relief from this procedure that persisted at their 9-month follow-up. The authors concluded that in patients with injection confirmed sacroiliac joint pain who respond to L4-L5 dorsal rami and S1-3 lateral branch blocks, radiofrequency denervation can be an effective treatment. Limitations of this study include the observational nature and small number of patients.

Characteristics of Excluded Studies

Among the studies failing to meet the strict criteria for this evaluation was a randomized, placebo-controlled study evaluating lateral branch radiofrequency denervation by Cohen et al (107). Except for dual blocks, the study meets all the criteria for randomized trials (117) and the reporting guidelines of CONSORT (118). This study was also the first to utilize cooled probe radiofrequency technology, which can increase the lesion size by a factor of 8 (31). The authors randomized 28 patients from amongst 90

potential candidates with predominantly axial low back pain to receive either cooled radiofrequency denervation from L4-S3 or sham lesioning. The main inclusion criterion was > 75% pain relief lasting at least 3 hours following a single intraarticular block performed with a 3 mL solution containing 2 mL of bupivacaine and 40 mg of depomethylprednisolone. Those patient's allocated to the placebo group who failed to obtain significant benefit were eligible to crossover to an open-label parallel group that received conventional radiofrequency denervation, 3 and 6 months after the procedure, 64% (n = 9) patients and 57% (n = 8) patients undergoing cooled radiofrequency lesioning experienced > 50% pain relief accompanied by significant functional improvement. In contrast, none of the sham-treated patients experienced significant improvement 3 months after the procedure. In the crossover treatment group (n = 11), 6 (55%) and 4 (36%) patients experienced a positive outcome 3 and 6 months post-procedure. However, one year after treatment, only 2 patients (14%) in the treatment group continued to demonstrate persistent pain relief. The authors concluded that these results furnished preliminary evidence that L4 and L5 primary dorsal rami and S1 to S3 lateral branch radiofrequency denervation may provide intermediate-term pain relief and functional benefit in well-selected patients with suspected sacroiliac joint pain. They also conceded that larger studies were needed to confirm these results and identify the optimal candidates and treatment parameters for this therapy.

This study provides strong evidence that response to radiofrequency denervation is superior to placebo. The limitations of the study include the small number of patients, the failure to exclude false-positive responders with a single uncontrolled sacroiliac joint block, the utilization of different types of radiofrequency technology, and the abridged outcome measures after 6 months.

Level of Evidence

Based on the available literature, evidence is unavailable for intraarticular sacroiliac joint injections for therapeutic purposes. For radiofrequency neurotomy, the indicated evidence is Level II-3 (limited). The recommendations based on Guyatt et al's (62) criteria are 2B/a weak recommendation for radiofrequency neurotomy for sacroiliac joint pain.

DISCUSSION

Based on this systematic review that included 5 studies evaluating the diagnostic accuracy of sacroiliac joint injections, the indicated evidence is Level II-2. For therapeutic interventions, there was no evidence supporting or refuting intraarticular injections. For radiofrequency neurotomy, the indicated evidence is Level II-3 or limited. The prevalence of sacroiliac joint pain is estimated to range between 10% and 38%, with the false-positive rate range between 20% and 54% for uncontrolled single blocks (4,9,10,71,72). This evaluation also determined that a combination of non-invasive provocative measures have limited ability to discriminate between sacroiliac joint pain and other lumbar spine disorders (level of evidence II-3). The results of this systematic review are similar to those from previous systematic reviews (1,2).

The inclusion criteria formulated for this review considered only those studies in which a double-diagnostic block paradigm was used to establish a painful sacroiliac joint. This was done for several reasons. First it was felt that using true placebo blocks for diagnostic purposes are generally considered to be unethical and impractical. Yet, without the use of double comparative blocks, one cannot reliably eliminate false-positive responders. The use of double-blocks to select patients for lumbar facet joint interventions is far more consistent than for SI joint therapies (4,31,44,48-50,119-124). Further confounding the use of double-blocks in selecting candidates for sacroiliac joint denervation are the uncertainties and vagaries surrounding the nerve supply (125). This has led some investigators to suggest that double sacroiliac joint injections are the most reliable means to select treatment candidates. Despite our advocacy for double-diagnostic selection criteria, the disparities in published studies indicate that this is not a universally accepted criterion. Furthermore, what little literature does exist on this topic suggests that using double blocks prior to radiofrequency denervation may not improve treatment outcomes (126).

A second controversial issue surrounds whether or not intraarticular injections are more advantageous than peri-articular injections. The ligamentous connections surrounding the SI joint are intricate and complex. Patients may exhibit either intra- or extra-articular SI pathology, but no reliable means exists to distinguish between the two. For therapeutic procedures, both randomized, controlled studies examining peri-articular corticosteroid injections in patients with

and without spondyloarthropathy demonstrated significant short-term benefit (83,91). Some patients may even present with both intra- and extraarticular pain generators. A retrospective review by Borowsky and Fagen (98) conducted in 120 patients found the combination of intra- and peri-articular injectate deposition provided superior analgesia than intraarticular injection alone.

The question of whether or not to screen patients for SI joint denervation using intra- or peri-articular injections is no less complicated. In fact, evidence may even support using peri-articular injections to select patients for lateral branch radiofrequency lesioning. In a study by Dreyfuss et al (127) conducted in asymptomatic volunteers, the authors found that multi-depth lateral branch injections blocked nociceptive input secondary to ligamentous probing (i.e., extraarticular stimulation) 70% of the time, compared to only 20% of the time during capsular distension (i.e., intraarticular stimulation). In the only clinical study that screened patients with extraarticular ligamentous injections, Yin et al (103) reported a 64% success rate 6-months after sacroiliac joint radiofrequency denervation.

Notwithstanding the contentious issues, based on the current level of evidence, the use of sacroiliac joint injections for diagnostic utilization is moderately supported. Similar to diagnostic utilization, the use of radiofrequency neurotomy remains in constant flux. However, considering that there is no other viable alternative to managing sacroiliac joint pain in patients refractory to corticosteroid injections, the judicious use of this technology in carefully selected patients appears warranted. But it is equally clear that further studies are needed to both refine the selection criteria and improve the technology.

The limitations of this systematic review include the scant literature available for analysis, the flawed methodology utilized in multiple evaluations, and the large scale discrepancies in techniques, outcome measures, and follow-up periods.

In summary, sacroiliac joint injections are safe and reasonable tools when used diagnostically, but therapeutic measures should be cautiously utilized based on strict selection criteria, in parallel with the physician's experience and technical abilities.

CONCLUSION

This systematic review lends moderate support for the use of diagnostic sacroiliac joint interven-

tions in chronic low back and/or lower extremity pain, whereas it provides limited evidence for radiofrequency neurotomy of sacroiliac joint nerve supply. Thus, a cautious approach must be utilized for diagnosis and even more cautious for therapeutic management, either with intraarticular injections or radiofrequency neurotomy based on individual patient circumstances and the clinician's experience and technical capabilities.

ACKNOWLEDGMENTS

The authors wish to thank the editorial board of *Pain Physician*, for review and criticism in improving the manuscript; Vidyasagar Pampati, MSc, statistician; Sekar Edem for assistance in search of literature; and Tonie M. Hatton and Diane E. Neihoff, transcriptionists (Pain Management Center of Paducah), for their assistance in preparation of this manuscript.

REFERENCES

- Hansen HC, McKenzie-Brown AM, Cohen SP, Swicegood JR, Colson JD, Manchikanti L. Sacroiliac joint interventions: A systematic review. *Pain Physician* 2007; 10:165-184.
- McKenzie-Brown A, Shah RV, Sehgal N, Everett CR. A systematic review of sacroiliac joint interventions. *Pain Physician* 2005; 8:115-125.
- Boswell MV, Trescot AM, Datta S, Schultz DM, Hansen HC, Abdi S, Sehgal N, Shah RV, Singh V, Benyamin RM, Patel VB, Buenaventura RM, Colson JD, Cordner HJ, Epter RS, Jasper JF, Dunbar EE, Atluri SL, Bowman RC, Deer TR, Swicegood JR, Staats PS, Smith HS, Burton AW, Klothe DS, Giordano J, Manchikanti L. Interventional techniques: Evidence-based practice guidelines in the management of chronic spinal pain. *Pain Physician* 2007; 10:7-111.
- Manchikanti L, Singh V, Pampati V, Dameron K, Barnhill R, Beyer C, Cash K. Evaluation of the relative contributions of various structures in chronic low back pain. *Pain Physician* 2001; 4:308-316.
- Cohen SP. Sacroiliac joint pain: A comprehensive review of anatomy, diagnosis and treatment. *Anesth Analg* 2005; 101:1440-1453.
- Foley BS, Buschbacher RM. Sacroiliac joint pain: Anatomy, biomechanics, diagnosis, and treatment. *Am J Phys Med Rehabil* 2006; 85:997-1006.
- Forst SL, Wheeler MT, Fortin JD, Vilensky JA. The sacroiliac joint: Anatomy, physiology, and clinical significance. *Pain Physician* 2006; 9:61-67.
- Zelle BA, Gruen GS, Brown S, George S. Sacroiliac joint dysfunction: Evaluation and management. *Clin J Pain* 2005; 21:446-455.
- Maigne JY, Aivakiklis A, Pfefer F. Results of sacroiliac joint double block and value of sacroiliac pain provocation test in 54 patients with low back pain. *Spine* 1996; 21:1889-1892.
- Irwin RW, Watson T, Minick RP, Ambrosius WT. Age, body mass index, and gender differences in sacroiliac joint pathology. *Am J Phys Med Rehabil* 2007; 86:37-44.
- Schwarzer AC, Aprill CN, Bogduk M. The sacroiliac joint in chronic low back pain. *Spine* 1995; 20:31-37.
- Maigne JY, Planchon CA. Sacroiliac joint pain after fusion. A study with anesthetic blocks. *Eur Spine J* 2005; 14:654-658.
- Merskey H, Bogduk N. Sacroiliac joint pain in Group XXVII: Sacral spinal or radicular pain syndromes. In *Classification of Chronic Pain: Descriptions of Chronic Pain Syndromes and Definition of Pain Terms*, 2nd ed. Task Force on Taxonomy of the International Association for the Study of Pain. IASP Press, Seattle, 1994, pp 190-191.
- Bogduk N. The sacroiliac joint. In *Clinical Anatomy of Lumbar Spine and Sacrum*, 4th edition. Churchill Livingstone, New York, 2005 pp 173-181.
- Murata Y, Takahashi K, Yamagata M, Takahashi Y, Shimada Y, Moriya H. Origin and pathway of sensory nerve fibers to the ventral and dorsal sides of the sacroiliac joint in rats. *J Orthop Res* 2001; 19:379-383.
- Fortin JD, Kissling RO, O'Connor BL, Vilensky JA. Sacroiliac joint innervation and pain. *Am J Orthop* 1999; 28:687-690.
- Grob KR, Neuhuber WL, Kissling RO. Innervation of the sacroiliac joint of the human. *Z Rheumatol* 1995; 54:117-122.
- Ikeda R. Innervation of the sacroiliac joint. Macroscopical and histological studies. *Nippon Ika Daigaku Zasshi* 1991; 58:587-596.
- Vilensky JA, O'Connor BL, Fortin JD, Merkel GJ, Jimenez AM, Scofield BA, Kleiner JB. Histologic analysis of neural elements in the human sacroiliac joint. *Spine* 2002; 27:1202-1207.
- Sakamoto N, Yamashita T, Takebayashi T, Sekine M, Ishii S. An electrophysiological study of mechanoreceptors in the sacroiliac joint and adjacent tissues. *Spine* 2001; 26:E468-E471.
- Solonen KA. The sacroiliac joint in the light of anatomical, roentgenological and clinical studies. *Acta Orthop Scand* 1957; 27:1-27.
- Minaki Y, Yamashita T, Ishii S. An electrophysiological study on the mechanoreceptors in the lumbar spine and adjacent tissues. *Neurol Orthop* 1996; 20:23-35.
- Manchikanti L, Boswell MV, Singh V, Hansen HC. Sacroiliac joint pain: Should physicians be blocking lateral branches, medial branches, dorsal rami, or ventral rami? (Letter to the Editor) *Reg Anesth Pain Med* 2003; 28:490-491.
- Szadek KM, Hoogland PV, Zuurmond WW, de Lange JJ, Perez RS. Nociceptive nerve fibers in the sacroiliac joint in humans. *Reg Anesth Pain Med* 2008; 33:36-43.
- Fortin JD, Dwyer AP, West S, Pier J. Sacroiliac joint: Pain referral maps upon applying a new injection/arthrography technique. Part I: Asymptomatic volunteers. *Spine* 1994; 19:1475-1482.
- Fortin JD, Aprill CN, Ponthieux B, Pier J. Sacroiliac joints: Pain referral maps upon applying a new injection/arthrography technique. Part II: Clinical evaluation. *Spine* 1994; 19:1483-1489.
- Slipman CW, Jackson HB, Lipetz JS, Chan KT, Lenrow D, Vresilovic EJ. Sacroiliac joint pain referral zones. *Arch Phys Med Rehabil* 2000; 81:334-338.
- Hancock MJ, Maher CG, Latimer J, Spindler MF, McAuley JH, Laslett M, Bogduk N. Systematic review of tests to identify

- the disc, SIJ or facet joint as the source of low back pain. *Eur Spine J* 2007; 16:1539-1550.
29. Buenaventura RM, Shah RV, Patel V, Benyamin R, Singh V. Systematic review of discography as a diagnostic test for spinal pain: An update. *Pain Physician* 2007; 10:147-164.
 30. Datta S, Everett CR, Trescot AM, Schultz DM, Adlaka R, Abdi S, Atluri SL, Smith HS, Shah RV. An updated systematic review of diagnostic utility of selective nerve root blocks. *Pain Physician* 2007; 10:113-128.
 31. Sehgal N, Dunbar EE, Shah RV, Colson JD. Systematic review of diagnostic utility of facet (zygapophysial) joint injections in chronic spinal pain: An update. *Pain Physician* 2007; 10:213-228.
 32. Wolfer L, Derby R, Lee JE, Lee SH. Systematic review of lumbar provocation discography in asymptomatic subjects with a meta-analysis of false-positive rates. *Pain Physician* 2008; 11:513-538.
 33. Manchikanti L, Singh V, Derby R, Schultz DM, Benyamin RM, Prager JP, Hirsch JA. Reassessment of evidence synthesis of occupational medicine practice guidelines for interventional pain management. *Pain Physician* 2008; 11:393-482.
 34. Saal JS. General principles of diagnostic testing as related to painful lumbar spine disorders. *Spine* 2002; 27:2538-2545.
 35. Hildebrandt J. Relevance of nerve blocks in treating and diagnosing low back pain – is the quality decisive? *Schmerz* 2001; 15:474-483.
 36. Szadek KM, van der Wurff P, van Tulder MW, Zuurmond WW, Perez RR. Diagnostic validity of criteria for sacroiliac joint pain: A systematic review. *J Pain* 2008; [Epub ahead of print].
 37. Song IH, Carrasco-Fernández J, Rudwaleit M, Sieper J. The diagnostic value of scintigraphy in assessing sacroiliitis in ankylosing spondylitis: A systematic literature research. *Ann Rheum Dis* 2008; 67:1535-1540.
 38. Rubinstein SM, van Tulder M. A best-evidence review of diagnostic procedures for neck and low-back pain. *Best Pract Res Clin Rheumatol* 2008; 22:471-482.
 39. Airaksinen O, Brox JI, Cedraschi C, Hildebrandt J, Klaber-Moffett J, Kovacs F, Mannion AF, Reis S, Staal JB, Ursin H, Zanoli G. Chapter 4: European guidelines for the management of chronic nonspecific low back pain. *Eur Spine J* 2006; 15:S192-S300.
 40. Manchikanti L, Singh V, Pampati V, Smith HS, Hirsch J. Analysis of growth of interventional techniques in managing chronic pain in Medicare population: A 10-year evaluation from 1997 to 2006. *Pain Physician* 2009; 12:9-34.
 41. Manchikanti L, Boswell MV. Interventional techniques in ambulatory surgical centers: A look at the new payment system. *Pain Physician* 2007; 10:627-650.
 42. Manchikanti L, Hirsch JA. Issues in Health Care: Interventional pain management at the crossroads. *Health Policy Update. Pain Physician* 2007; 10:261-284.
 43. West S, King V, Carey TS, Lohr KN, McKeoy N, Sutton SF, Lux L. *Systems to Rate the Strength of Scientific Evidence*. Evidence Report, Technology Assessment No. 47. AHRQ Publication No. 02-E016. Rockville, MD: Agency for Health care Research and Quality, 2002. www.thecre.com/pdf/ahrq-system-strength.pdf
 44. Atluri S, Datta S, Falco FJE, Lee M. Systematic review of diagnostic utility and therapeutic effectiveness of thoracic facet joint interventions. *Pain Physician* 2008; 11:611-629.
 45. Singh V, Manchikanti L, Shah RV, Dunbar EE, Glaser SE. Systematic review of thoracic discography as a diagnostic test for chronic spinal pain. *Pain Physician* 2008; 11:631-642.
 46. Manchikanti L, Dunbar EE, Wargo BW, Shah RV, Derby R, Cohen SP. Systematic Review of Cervical Discography as a Diagnostic Test for Chronic Spinal Pain. *Pain Physician* 2009; 12:305-321.
 47. Wolfer L, Derby R, Lee JE, Lee SH. Systematic review of lumbar provocation discography in asymptomatic subjects with a meta-analysis of false-positive rates. *Pain Physician* 2008; 11:513-538.
 48. Datta S, Lee M, Falco FJE, Bryce DA, Hayek SM. Systematic Assessment of Diagnostic Accuracy and Therapeutic Utility of Lumbar Facet Joint Interventions. *Pain Physician* 2009; 12:437-460.
 49. Falco FJE, Erhart S, Wargo BW, Bryce DA, Atluri S, Datta S, Hayek SM. Systematic Review of Diagnostic Utility and Therapeutic Effectiveness of Cervical Facet Joint Interventions. *Pain Physician* 2009; 12:323-344.
 50. Smith HS, Chopra P, Patel VB, Frey ME, Rastogi R. Systematic review on the role of sedation in diagnostic spinal interventional techniques. *Pain Physician* 2009; 12:195-206.
 51. Berg AO, Allan JD. Introducing the third U.S. Preventive Services Task Force. *Am J Prev Med* 2001; 20:S3-S4.
 52. Koes BW, Scholten RJ, Mens JM, Bouter LM. Efficacy of epidural steroid injections for low-back pain and sciatica: A systematic review of randomized clinical trials. *Pain* 1995; 63:279-288.
 53. Helm S, Hayek S, Benyamin RM, Manchikanti L. Systematic review of effectiveness of thermal annular procedures in treating discogenic low back pain. *Pain Physician* 2009; 12:207-232.
 54. Conn A, Buenaventura RM, Datta S, Abdi S, Diwan S. Systematic review of caudal epidural injections in the management of chronic low back pain. *Pain Physician* 2009; 12:109-135.
 55. Benyamin RM, Singh V, Parr AT, Conn A, Diwan S, Abdi S. Systematic review of the effectiveness of cervical epidurals in the management of chronic neck pain. *Pain Physician* 2009; 12:137-157.
 56. Parr AT, Diwan S, Abdi S. Lumbar interlaminar epidural injections in managing chronic low back and lower extremity pain: A systematic review. *Pain Physician* 2009; 12:163-188.
 57. Buenaventura RM, Datta S, Abdi S, Smith HS. Systematic review of therapeutic lumbar transforaminal epidural steroid injections. *Pain Physician* 2009; 12:233-251.
 58. Epter RS, Helm S, Hayek SM, Benyamin RM, Smith HS, Abdi S. Systematic Review of Percutaneous Adhesiolysis and Management of Chronic Low Back Pain in Post Lumbar Surgery Syndrome. *Pain Physician* 2009; 12:361-378.
 59. Hayek SM, Helm S, Benyamin RM, Singh V, Bryce DA, Smith HS. Effectiveness of Spinal Endoscopic Adhesiolysis in Post Lumbar Surgery Syndrome: A Systematic Review. *Pain Physician* 2009; 12:419-435.
 60. Frey ME, Manchikanti L, Benyamin RM, Schultz DM, Smith HS, Cohen SP. Spinal Cord Stimulation for Patients with Failed Back Surgery Syndrome: A Systematic Review. *Pain Physician* 2009; 12:379-397.
 61. Patel VB, Manchikanti L, Singh V, Schultz DM, Hayek SM, Smith HS. Systematic Review of Intrathecal Infusion Systems for Long-Term Management of Chronic

- Non-Cancer Pain. *Pain Physician* 2009; 12:345-360.
62. Guyatt G, Gutterman D, Baumann MH, Addrizzo-Harris D, Hylek EM, Phillips B, Raskob G, Lewis SZ, Schünemann H. Grading strength of recommendations and quality of evidence in clinical guidelines. Report from an American College of Chest Physicians task force. *Chest* 2006; 129:174-181.
 63. Dreyfuss P, Michaelsen M, Pauza K, McLarty J, Bogduk N. The value of medical history and physical examination in diagnosing sacroiliac joint pain. *Spine* 1996; 21:2594-2602.
 64. Laslett M, Aprill CN, McDonald B, Young SB. Diagnosis of sacroiliac joint pain: A validity of individual provocation tests and composites of tests. *Man Ther* 2005; 10:207-218.
 65. Young S, Aprill CN, Laslett M. Correlation of clinical examination characteristics with three sources of chronic low back pain. *Spine J* 2003; 3:460-465.
 66. Maigne JY, Boulahdour H, Chatellier G. Value of quantitative radionuclide bone scanning in the diagnosis of sacroiliac joint syndrome in 32 patients with low back pain. *Eur Spine J* 1998; 7:328-331.
 67. Slipman CW, Sterenfeld EB, Chou LH, Herzog R, Vresilovic E. The value of radionuclide imaging in the diagnosis of sacroiliac joint syndrome. *Spine* 1996; 21:2251-2254.
 68. Broadhurst NA, Bond MJ. Pain provocation tests for the assessment of sacroiliac joint dysfunction. *J Spin Disord* 1998; 11:341-345.
 69. Bogduk N. Pain provocation tests for the assessment of sacroiliac joint dysfunction. *J Spinal Disord* 1999; 12:357-358.
 70. Pang WW, Mok MS, Lin ML, Chang DP, Hwang MH. Application of spinal pain mapping in the diagnosis of low back pain - Analysis of 104 cases. *Acta Anaesthesiol Sin* 1998; 36:71-74.
 71. Laslett M, Young SB, Aprill CN, McDonald B. Diagnosing painful sacroiliac joints: A validity study of a McKenzie evaluation and sacroiliac provocation tests. *Aust J Physiother* 2003; 49:89-97.
 72. van der Wurff P, Buijs EJ, Groen GJ. A multitest regimen of pain provocation tests as an aid to reduce unnecessary minimally invasive sacroiliac joint procedures. *Arch Phys Med Rehabil* 2006; 87:10-14.
 73. van der Wurff P, Buijs EJ, Groen GJ. Intensity mapping of pain referral areas in sacroiliac joint pain patients. *J Manipulative Physiol Ther* 2006; 29:190-195.
 74. Berthelot JM, Labat JJ, Le Goff B, Gouin F, Maugars Y. Provocative sacroiliac joint maneuvers and sacroiliac joint block are unreliable for diagnosing sacroiliac joint pain. *Joint Bone Spine* 2006; 73:17-23.
 75. Simpson R, Gemmell H. Accuracy of spinal orthopaedic tests: A systemic review. *Chiropr Osteopat* 2006; 14:26.
 76. Leboeuf C. The sensitivity and specificity of seven lumbo-pelvic orthopedic tests and the arm-fossa test. *J Manip Physiol Ther* 1990; 13:138-143.
 77. Jung JH, Kim HI, Shin DA, Shin DG, Lee JO, Kim HJ, Chung JH. Usefulness of pain distribution pattern assessment in decision-making for the patients with lumbar zygapophyseal and sacroiliac joint arthropathy. *J Korean Med Sci* 2007; 22:1048-1054.
 78. Puhakka KB, Jurik AG, Schiottz-Christensen B, Hansen GV, Egund N, Christiansen JV, Stengaard-Pedersen K. MRI abnormalities of sacroiliac joints in early spondyloarthropathy: A 1-year follow-up study. *Scand J Rheumatol* 2004; 33:332-338.
 79. Puhakka KB, Melson F, Jurik AG, Boel LW, Vesterby A, Egund N. MR imaging of the normal sacroiliac joint with correlation to histology. *Skeletal Radiol* 2004; 33:15-28.
 80. Vogler JB 3rd, Brown WH, Helms CA, Genant HK. The normal sacroiliac joint: A CT study of asymptomatic patients. *Radiology* 1984; 151:433-437.
 81. Elgafy H, Semaan HB, Ebraheim NA, Coombs RJ. Computed tomography findings in patients with sacroiliac pain. *Clin Orthop Relat Res* 2001; 382:112-118.
 82. Dussault RG, Kaplan PA, Anderson MW. Fluoroscopy-guided sacroiliac joint injections. *Radiology* 2000; 214:273-277.
 83. Luukkainen RK, Wennerstrand PV, Kautiainen HH, Sanila MT, Asikainen EL. Efficacy of periarticular corticosteroid treatment of the sacroiliac joint in non-spondylarthropathic patients with chronic low back pain in the region of the sacroiliac joint. *Clin Exp Rheumatol* 2002; 20:52-54.
 84. Maugars Y, Mathis C, Berthelot JM, Charlier C, Prost A. Assessment of the efficacy of sacroiliac corticosteroid injections spondyloarthropathies: A double blind study. *Br J Rheumatol* 1996; 35:767-770.
 85. Karabacakoglu A, Karakose S, Ozerbil OM, Odev K. Fluoroscopy-guided intraarticular corticosteroid injection into the sacroiliac joints in patients with ankylosing spondylitis. *Acta Radiol* 2002; 43:425-427.
 86. Fischer T, Biedermann T, Hermann KG, Diekmann F, Braun J, Hamm B, Bollow M. Sacroiliitis in children with spondyloarthropathy: Therapeutic effect of CT-guided intraarticular corticosteroid injection. *Rofa* 2003; 175:814-821.
 87. Hanly JG, Mitchell M, MacMillan L, Mosher D, Sutton E. Efficacy of sacroiliac corticosteroid injections in patients with inflammatory spondyloarthropathy: Results of a 5 month controlled study. *J Rheum* 2000; 27:719-722.
 88. Maugars Y, Mathis C, Vilon P, Prost A. Corticosteroid injection of the sacroiliac joint in patients with seronegative spondylarthropathy. *Arthritis Rheum* 1992; 35:564-568.
 89. Bollow M, Braun J, Taupitz M, Haberle J, Reibhauer BH, Paris S, Mutze S, Seyrekbasan F, Wolf KJ, Hamm B. CT-guided intraarticular corticosteroid injection into the sacroiliac joints in patients with spondyloarthropathy: Indication and follow-up with contrast-enhanced MRI. *J Comput Assist Tomogr* 1996; 20:512-521.
 90. Braun J, Bollow M, Seyrekbasan F, Haberle HJ, Eggens U, Mertz A, Distler A, Sieper J. Computed tomography guided corticosteroid injection of the sacroiliac joint in patients with spondyloarthropathy with sacroiliitis: Clinical outcome and follow-up by dynamic magnetic resonance imaging. *J Rheumatol* 1996; 23:659-664.
 91. Luukkainen R, Nissila M, Asikainen E, Sanila M, Lehtinen K, Alanaatu A, Kautiainen H. Periarticular corticosteroid treatment of the sacroiliac joint in patients with seronegative spondyloarthropathy. *Clin Exp Rheumatol* 1999; 17:88-90.
 92. Gunaydin I, Pereira PL, Daikeler T, Mohren M, Trubenbach J, Schick F, Kanz L, Kotter I. Magnetic resonance imaging guided corticosteroid injection of the sacroiliac joints in patients with therapy resistant spondyloarthropathy: A pilot study. *J Rheumatol* 2000; 27:424-428.
 93. Pereira PL, Gunaydin I, Duda SH, Tru-

- benbach J, Remy CT, Kotter I, Kastler B, Claussen CD. Corticosteroid injections of the sacroiliac joint during magnetic resonance: Preliminary results [in French]. *J Radiol* 2000; 81:223-226.
94. Pereira PL, Gunaydin I, Trubenbach J, Dammann F, Remy CT, Kotter I, Schick F, Koenig CW, Claussen CD. Interventional MR imaging for injection of sacroiliac joints in patients with sacroiliitis. *AJR Am J Roentgenol* 2000; 175:265-266.
 95. Ojala R, Klemola R, Karppinen J, Sequeiros RB, Tervonen O. Sacro-iliac joint arthrography in low back pain: Feasibility of MRI guidance. *Eur J Radiol* 2001; 40:236-239.
 96. Slipman CW, Lipetz JS, Plastaras CT, Jackson HB, Vresilovic EJ, Lenrow DA, Braverman DL. Fluoroscopically guided therapeutic sacroiliac joint injections for sacroiliac joint syndrome. *Am J Phys Med Rehabil* 2001; 80:425-432.
 97. Chakraverty R, Dias R. Audit of conservative management of chronic low back pain in a secondary care setting - Part I: Facet joint and sacroiliac joint interventions. *Acupunct Med* 2004; 22:207-213.
 98. Borowsky CD, Fagen G. Sources of sacroiliac region pain: Insights gained from a study comparing standard intraarticular injection with a technique combining intra- and peri-articular injection. *Arch Phys Med Rehabil* 2008; 89:2048-2056.
 99. Murakami E, Tanaka Y, Aizawa T, Ishizuka M, Kokubun S. Effect of periarticular and intraarticular lidocaine injections for sacroiliac joint pain: Prospective comparative study. *J Orthop Sci* 2007; 12:274-280.
 100. Ferrante FM, King LF, Roche EA, Kim PS, Aranda M, Delaney LR, Mardini IA, Mannes AJ. Radiofrequency sacroiliac joint denervation for sacroiliac syndrome. *Reg Anesth Pain Med* 2001; 26:137-142.
 101. Gevargez A, Groenemeyer D, Schirp S, Braun M. CT-guided percutaneous radiofrequency denervation of the sacroiliac joint. *Eur Radiol* 2002; 12:1360-1365.
 102. Cohen SP, Abdi S. Lateral branch blocks as a treatment for sacroiliac joint pain: A pilot study. *Reg Anesth Pain Med* 2003; 28:113-119.
 103. Yin W, Willard F, Carreiro J, Dreyfuss P. Sensory stimulation-guided sacroiliac joint radiofrequency neurotomy: Technique based on neuroanatomy of the dorsal sacral plexus. *Spine* 2003; 28:2419-2425.
 104. Buijs EJ, Kamphuis ET, Groen GJ. Radiofrequency treatment of sacroiliac joint-related pain aimed at the first three sacral dorsal rami: A minimal approach. *Pain Clinic* 2004; 16:139-146.
 105. Burnham RS, Yasui Y. An alternate method of radiofrequency neurotomy of the sacroiliac joint: A pilot study of the effect of pain, function, and satisfaction. *Reg Anesth Pain Med* 2007; 32:12-19.
 106. Vallejo R, Benyamin RM, Kramer J, Stanton G, Joseph NJ. Pulsed radiofrequency denervation for the treatment of sacroiliac joint syndrome. *Pain Med* 2006; 7:429-434.
 107. Cohen SP, Hurley RW, Buckenmaier CC 3rd, Kurihara C, Morlando B, Dragovich A. Randomized placebo-controlled study evaluating lateral branch radiofrequency denervation for sacroiliac joint pain. *Anesthesiology* 2008; 109:279-288.
 108. Kapural L, Nageeb F, Kapural M, Cata JP, Narouze S, Mekhail N. Cooled radiofrequency system for the treatment of chronic pain from sacroiliitis: The first case-series. *Pain Pract* 2008; 8:348-354.
 109. Harmon D, O'Sullivan M. Ultrasound-guided sacroiliac joint injection technique. *Pain Physician* 2008; 11:543-547.
 110. Manchikanti L, Singh V, Falco FJE, Cash KA, Pampati V. Effectiveness of thoracic medial branch blocks in managing chronic pain: A preliminary report of a randomized, double-blind controlled trial: Clinical Trial NCT00355706. *Pain Physician* 2008; 11:491-504.
 111. Manchikanti L, Singh V, Falco FJ, Cash KA, Fellows B. Cervical medial branch blocks for chronic cervical facet joint pain: A randomized double-blind, controlled trial with one-year follow-up. *Spine* 2008; 33:1813-1820.
 112. Manchikanti L, Singh V, Falco FJ, Cash KA, Pampati V. Lumbar facet joint nerve blocks in managing chronic facet joint pain: One-year follow-up of a randomized, double-blind controlled trial: Clinical Trial NCT00355914. *Pain Physician* 2008; 11:121-132.
 113. Manchikanti L, Cash KA, McManus CD, Pampati V, Smith HS. Preliminary results of randomized, equivalence trial of fluoroscopic caudal epidural injections in managing chronic low back pain: Part 1. Discogenic pain without disc herniation or radiculitis. *Pain Physician* 2008; 11:785-800.
 114. Manchikanti L, Singh V, Cash KA, Pampati V, Damron KS, Boswell MV. Preliminary results of randomized, equivalence trial of fluoroscopic caudal epidural injections in managing chronic low back pain: Part 2. Disc herniation and radiculitis. *Pain Physician* 2008; 11:801-815.
 115. Manchikanti L, Singh V, Cash KA, Pampati V, Datta S. Preliminary results of randomized, equivalence trial of fluoroscopic caudal epidural injections in managing chronic low back pain: Part 3. Post surgery syndrome. *Pain Physician* 2008; 11:817-831.
 116. Manchikanti L, Cash KA, McManus CD, Pampati V, Abdi S. Preliminary results of randomized, equivalence trial of fluoroscopic caudal epidural injections in managing chronic low back pain: Part 4. Spinal stenosis. *Pain Physician* 2008; 11:833-848.
 117. Manchikanti L, Hirsch JA, Smith HS. Evidence-based medicine, systematic reviews, and guidelines in interventional pain management: Part 2: Randomized controlled trials. *Pain Physician* 2008; 11:717-773.
 118. Altman DG, Schulz KF, Moher D, Egger M, Davidoff F, Elbourne D, Gotzsche PC, Lang T; CONSORT GROUP (Consolidated Standards of Reporting Trials). The revised CONSORT statement for reporting randomized trials: Explanation and elaboration. *Ann Intern Med* 2001; 134:663-694.
 119. Boswell MV, Colson JD, Sehgal N, Dunbar E, Eptner R. A systematic review of therapeutic facet joint interventions in chronic spinal pain. *Pain Physician* 2007; 10:229-253.
 120. Manchikanti L, Boswell MV, Singh V, Pampati V, Damron KS, Beyer CD. Prevalence of facet joint pain in chronic spinal pain of cervical, thoracic, and lumbar regions. *BMC Musculoskelet Disord* 2004; 5:15.
 121. Manchukonda R, Manchikanti KN, Cash KA, Pampati V, Manchikanti L. Facet joint pain in chronic spinal pain: An evaluation of prevalence and false-positive rate of diagnostic blocks. *J Spinal Disord Tech* 2007; 20:539-545.
 122. Manchikanti L, Cash KA, Pampati V, Fellows B. Influence of psychological variables on the diagnosis of facet joint involvement in chronic spinal pain. *Pain Physician* 2008; 11:145-160.

123. Manchikanti L, Manchikanti K, Cash KA, Singh V, Giordano J. Age-related prevalence of facet joint involvement in chronic neck and low back pain. *Pain Physician* 2008; 11:67-75.
124. Manchikanti L, Boswell MV, Manchukonda R, Cash KA, Giordano. Influence of prior opioid exposure on diagnostic facet joint nerve blocks. *J Opioid Manage* 2008; 4:351-360.
125. Cohen SP. Epidemics, evolution and SI joint pain. *Reg Anesth Pain Med* 2007; 32:3-6.
126. Cohen SP, Strassels S, Kurihara C, Crooks MT, Bleckner LL, Forsythe A, Marcuson M. Outcome predictors for sacroiliac (lateral branch) radiofrequency denervation. *Reg Anesth Pain Med* 2009; in press.
127. Dreyfuss P, Henning T, Malladi R, Goldstein B, Bogduk N. The ability of multi-site, multi-depth sacral lateral branch blocks to anesthetize the sacroiliac joint complex: A physiologic study. Presented at International Spinal Intervention Society 16th Annual Meeting, Las Vegas, NV, July 23-26, 2008.