

## Observational Study



# Visit Frequency and Outcomes for Patients Using Ongoing Chiropractic Care for Chronic Low-Back and Neck Pain: An Observational Longitudinal Study

Patricia M. Herman, PhD<sup>1</sup>, Sarah E. Edgington, MA<sup>1</sup>, Melony E. Sorbero, PhD<sup>1</sup>,  
Eric L. Hurwitz, PhD<sup>2</sup>, Christine M. Goertz, PhD<sup>3</sup>, and Ian D. Coulter, PhD<sup>1</sup>

From: <sup>1</sup>RAND Corporation, Santa Monica, CA; <sup>2</sup>Office of Public Health Studies, University of Hawaii, Honolulu, HI; <sup>3</sup>Duke University, Durham, NC

Address Correspondence:  
Patricia M. Herman, PhD  
Senior Behavioral/Social Scientist,  
RAND Corporation  
1776 Main Street, PO Box 2138  
Santa Monica, CA 90407-2138  
E-mail: pherman@rand.org

Disclaimer: The authors wish to disclose and thank the sponsor of the study. Our work on the project was funded by 2 grants (one from the National Center for Complementary and Integrative Health and one from the NCMIC Foundation) paid to RAND Corporation. The data used in this study were collected under a cooperative agreement (U19) from the National Center for Complementary and Integrative Health (NCCIH) Grant No. 1U19AT007912-01. The analyses in this study were funded by a grant from the National Chiropractic Mutual Insurance Company (NCMIC) Foundation. Our funders had no influence or interference after the protocol was originally proposed. The financial sponsors of this work had no role in the design and conduct of the study or the collection, management, analysis, and interpretation of the data. The sponsors also did not have a role in the preparation or review of the manuscript or the decision to submit.

Conflict of interest: Each author certifies that he or she, or a member of his or her immediate family, has no commercial association (i.e., consultancies, stock ownership, equity interest, patent/licensing arrangements, etc.) that might pose a conflict of interest in connection with the submitted manuscript.

Manuscript received: 05-26-2020  
Revised manuscript received: 07-09-2020  
Accepted for publication: 07-27-2020

Free full manuscript:  
[www.painphysicianjournal.com](http://www.painphysicianjournal.com)

**Background:** Chronic spinal pain is prevalent and long-lasting. Although provider-based nonpharmacologic therapies, such as chiropractic care, have been recommended, healthcare and coverage policies provide little guidance or evidence regarding long-term use of this care.

**Objective:** To determine the relationships between visit frequency and outcomes for patients using ongoing chiropractic care for chronic spinal pain.

**Study Design:** Observational 3-month longitudinal study.

**Setting:** Data collected from patients of 124 chiropractic clinics in 6 United States regions.

**Methods:** We examined the impact of visit frequency and patient characteristics on pain (pain 0-10 numeric rating scale) and functional outcomes (Oswestry Disability Index [ODI] for low-back pain and Neck Disability Index [NDI] for neck pain, both 0-100 scale) using hierarchical linear modeling (HLM) in a large national sample of chiropractic patients with chronic low back pain (CLBP) and/or chronic neck pain (CNP). This study was approved by the RAND Human Subjects Protection Committee and registered under ClinicalTrials.gov Identifier: NCT03162952.

**Results:** One thousand, three hundred, sixty-two patients with CLBP and 1,214 with CNP were included in a series of HLM models. Unconditional (time-only) models showed patients on average had mild pain and function, and significant, but slight improvements in these over the 3-month observation period: back and neck pain decreased by 0.40 and 0.44 points, respectively; function improved by 2.7 (ODI) and 3.0 points (NDI) (all  $P < 0.001$ ). Adding chiropractic visit frequency to the models revealed that those with worse baseline pain and function used more visits, but only visits more than once per week for those with CLBP were associated with significantly better improvement. These relationships remained when other types of visits and baseline patient characteristics were included.

**Limitations:** This is an observational study based on self-reported data from a sample representative of chiropractic patients, but not all patients with CLBP or CNP.

**Conclusions:** This 3-month window on chiropractic patients with CLBP and/or CNP revealed that they were improving, although slowly; may have reached maximum therapeutic improvement; and are possibly successfully managing their chronic pain using a variety of chiropractic visit frequencies. These results may inform payers when building coverage policies for ongoing chiropractic care for patients with chronic pain.

**Key words:** Chronic low back pain, chronic neck pain, spinal pain, physical function, hierarchical linear modeling, healthcare utilization, chiropractic visits, insurance coverage

**Pain Physician 2021: 24:E61-E74**

**A**lthough chronic pain affects over 40 percent of adults in the United States (US), little information is available on the management of that pain with ongoing provider-based care (1). Chronic spinal pain is one of the most common types of chronic pain (1,2). It is associated with a substantial burden to patients, the healthcare system, and employers (3-10). Although the use of medications (including opioids) is most common, provider-based nonpharmacologic therapies are now recommended in guidelines as first-line therapies for chronic spinal pain (5,11-16).

According to NIH Medline Plus, "chronic pain usually cannot be cured, but it can be managed" (17). Many turn to sustained medication use for this purpose. However, this approach has risks that may outweigh the benefits (18,19). We need information on long-term pain management for chronic low back pain (CLBP) and chronic neck pain (CNP) that includes the use of provider-based nonpharmacologic therapies. At present there is little guidance and sparse evidence available for this use (20-26).

Chiropractors, osteopaths, and physical therapists are the practitioners most likely to use spinal manipulation, one recommended provider-based nonpharmacologic therapy (27). About 30 to 60 percent of patients in the US with spinal pain have seen a chiropractor, and over 80 percent of chiropractic patients receive spinal manipulation for their back and neck pain (5,11,27,28). Most chiropractic patients have chronic pain and many are under long-term chiropractic care and very satisfied with this care (27,29-34). Therefore, an examination of visit frequencies, pain, and functional outcomes in patients using ongoing chiropractic care could be useful to understanding the use of provider-based nonpharmacologic therapies for pain management.

The available recommendations for provider-based care tend to give a frequency and duration of treatment (e.g., 10 treatments over 8 weeks) and the timing for reassessment before continuing the care plan (20-26). These treatment guidelines also refer to concepts like Maximum Therapeutic Improvement (MTI) (21,23,25,26): the point at which a patient's condition has plateaued and is unlikely to improve further (21). The guidelines all acknowledge that care beyond the point of MTI (i.e., chronic pain management or support care)—might be needed under certain conditions (e.g., if symptoms worsen after a therapeutic withdrawal of treatment) (21,22). One guideline suggests that pain or function must worsen by the minimal

clinically important change for more than 24 hours to justify ongoing care (21). However, although duration of care guidelines for chronic pain patients not yet at their MTI seem to be loosely based on treatment frequency and duration used in trials, little guidance and no evidence is offered for care after MTI is achieved. This lack of information and support of ongoing pain management has been cited as one barrier to the use of recommended provider-based nonpharmacologic therapies for chronic spinal pain (35).

This study used a longitudinal dataset, gathered over 3 months, from a large US sample of patients with CLBP and/or CNP who were using chiropractic care (29). While this sample may not be representative of all patients with CLBP and CNP, it is representative of chiropractic patients with CLBP and CNP (29,36), including pain and function levels seen in trials (37,38). We know from previous analyses of this sample that on average these patients have been in pain for 14 years and using chiropractic care for 11 years, and that 70 percent reported their treatment goal as pain management, not to cure (29,39). Their stated willingness-to-pay for pain reduction indicated that what they value is the maintenance of their current pain levels (40). On average they utilized 2.3 chiropractic visits per month, but this varied by the characteristics of the patient (more visits with worse function, just starting care, and with CLBP and insurance coverage) and their treating chiropractor (more visits when chiropractor saw more patients per day or had fewer years of experience) (36).

In this study, we examine relationships between patients' pain and functional outcomes over the 3-month study period, and their chiropractic visit frequency, visits to other types of providers, and other characteristics.

## **METHODS**

Our longitudinal, observational data were collected prospectively via online questionnaires every 2 weeks over 3 months from a large sample of US chiropractic patients under treatment for CLBP and/or CNP. The overall project, within which these data were collected, is described in detail elsewhere, including data collection methods, patient sample characteristics, and clinic and chiropractor characteristics (29,36,41-43). In brief, the sample was selected using multistage systematic stratified sampling over 4 levels: regions/states, metropolitan areas, chiropractors/clinics, and patients, and data were collected between October 2016 through January 2017. The regions and metropolitan areas were: San Diego, California; Tampa, Florida; Min-

neapolis, Minnesota; Seneca Falls/Upstate, New York; Portland, Oregon; and Dallas, Texas.

The goal was to recruit 20 chiropractors/clinics per region and 7 CLBP and 7 CNP cases per clinic. Each participating clinic received an iPad containing a short prescreening questionnaire and staff were trained to offer this questionnaire to every patient who visited the clinic during a 4-week period. The questionnaire established initial inclusion criteria (i.e., > 21 years of age, English-proficient, no current personal injury or workers compensation litigation/claims, have back or neck pain). Those who met these criteria and provided an email address were sent a link to a longer online screening questionnaire to determine whether they had CLBP and/or CNP (i.e., pain for at least 3 months before seeing the chiropractor and/or self-report of chronicity). Patients who met these criteria provided informed consent, answered additional questions, and then received 7 additional online questionnaires: baseline, 5 short every-2-week follow-ups, and endline at 3 months. Patients were incentivized with online gift cards for every step of participation and those who completed all questionnaires received a total of \$200. This study uses a subset of the data collected.

### Outcome Measures

The outcome and visit data were gathered every 2 weeks, over 3 months and exact weeks since baseline (based on actual date of data entry) was used as our time variable in the models. At each data collection point, all patients reported their pain levels (pain numeric rating scale [NRS]), their function using the NDI for those with CNP, and the ODI for those with CLBP (44-46). These measures are considered valid and reliable and were scored that higher values indicated worse outcomes (pain NRS (47-51); NDI (52-55); ODI (56-58)).

Our primary explanatory variable was chiropractic visit frequency, but we also tested the impact of other types of visits to other complementary therapy (CT) providers (mostly massage) and to medical providers (mostly general practitioners). Average frequency for each type of visit was categorized as: more than weekly, weekly up to biweekly, biweekly up to monthly, and often than monthly and less often than monthly. Not all patients had nonchiropractic visits and few used these more than biweekly, so for nonchiropractic visits we combined the first 2 categories and added a none category. Variables for clinic (chiropractor) and region (state and metropolitan area) were used to determine

whether there were differences in baseline symptoms or symptom change by chiropractor or region.

Our final models also included a number of variables identified by others as reasonable indicators of the need for ongoing care or shown to be predictive of outcomes in CLBP and CNP populations (21,38,59-72). These include pain characteristics (whether they have both CLBP and CNP, years with pain, reinjury tendency with heavy labor, or previous unsuccessful surgery), use of medications (over-the-counter and prescription pain medications, including narcotics), self-care (exercise), stage of care (first month or near end of care), pain beliefs (believe pain is chronic, pain level that would occur if they didn't see chiropractor, unsafe to be physically active/fear avoidance), psychological measures (pain management subscale of the Chronic Pain Self-Efficacy Scale, 2 items from the Credibility/Expectancy Questionnaire relating to expected treatment success and expected pain improvement, an item about worry whether pain will end, the 4-item PROMIS-29 depression scale (scores > 52.5), 3-item affective distress domain of the Multidimensional Pain Inventory, 3 items from Helplessness subscale of the Pain Catastrophizing Scale, and demographics (age, gender, education) (73-77). Each was chosen for the analysis a priori.

### Analysis

The goal of our analysis was to examine whether visit frequencies and patient characteristics were associated with patients' baseline pain and function, and with changes in these outcomes (i.e., more improvement) over the 3-month study period. Because we had up to 7 data points for each patient and patients were clustered within clinics and clinics were clustered within regions, we used hierarchical linear modeling (HLM) that corrects for error structure violations (e.g., non-independent errors), and optimizes estimation in the presence of missing data.

We first estimated unconditional (time-only) HLM models for each outcome to examine general trends in outcomes and to determine whether baseline patient symptoms or improvement over time varied significantly by chiropractor/clinic, and/or by region.

We then added frequency of chiropractic visit categories to unconditional models that appropriately clustered by patients, clinic, and/or region to examine the relationship between chiropractic visit frequency and outcomes. Next, we added other types of visits and then all explanatory variables in the full models.

Because of the large number of variables consid-

ered, at each step we used deviance statistics (measures of model fit based on log restricted-likelihood values of nested models) to separately test whether each block of variables was worth keeping—i.e., added statistically significant ( $P < 0.05$ ) explanatory power. We separately examined the power of each set of variables to explain baseline outcomes (main effects) and to explain changes in outcomes over the 3-month period (interactions with time/weeks).

Means and frequencies for all variables were compared across pain groups (CLBP only, both CLBP and CNP, and CNP only) using t-tests and  $\chi^2$ . All analyses were performed in Stata 16.0 (StataCorp, College Station, TX). This study was approved by the RAND Human Subjects Protection Committee.

## RESULTS

Of the 2,024 patients who completed the baseline survey, 1,708 had nonspecific CLBP or CNP, and 1,665 (97.5%) of those had sufficient data to be included in at least one of our HLM models (29) (Fig. 1). Table 1 shows the values of each variable considered by chronic pain type. In our sample, it was most common to have both CLBP and CNP, and these participants had more back pain, had their pain longer, were more likely to have had unsuccessful back surgery, were less likely to be a new patient or to be near to ending care, and had lower pain management self-efficacy and more worry about their pain, depression, affective distress, and catastrophizing than those with CLBP or CNP only. On average, over the 3-month period, patients in the sample




Fig. 1. Flow chart of patients into the study

## Chiropractic Visit Frequency Impact on Outcomes

Table 1. Outcomes and patient characteristics predictors by type of chronic pain mean (SD) unless otherwise noted.

Label	CLBP (n = 451)	CLBP & CNP (n = 911)	CNP (n = 303)	Totals (n = 1665)
Rating of Low Back Pain (0-10)**	3.3 (1.9)	3.6 (2.1)		3.5 (2.0)
Rating of Neck Pain (0-10)		3.8 (2.2)	3.5 (2.1)	3.7 (2.1)
Oswestry Disability Index Score (0-100)	18.5 (11.8)	19.8 (12.5)		19.4 (12.3)
Neck Disability Index Score (0-100)		21.8 (13.2)	20.7 (10.9)	21.5 (12.7)
# Chiropractic visits over 3 months*	5.5 (6.1)	5.7 (5.7)	6.3 (5.9)	6.0 (5.9)
Chiropractic visit frequency (categorized)**				
Monthly and less than monthly	190 (42.1%)	295 (32.4%)	107 (35.3%)	592 (35.6%)
Biweekly up to monthly	100 (22.2%)	251 (27.6%)	82 (27.1%)	433 (26.0%)
Weekly up to biweekly	84 (18.6%)	217 (23.8%)	62 (20.5%)	363 (21.8%)
More than weekly	48 (10.6%)	114 (12.5%)	31 (10.2%)	193 (11.6%)
Unknown	29 (6.4%)	34 (3.7%)	21 (6.9%)	84 (5.0%)
# Nonchiropractic complementary therapy (CT) visits over 3 months	1.8 (4.7)	2.2 (4.2)	2.2 (4.8)	2.1 (4.7)
Other CT visit frequency (categorized)**				
None	269 (59.6%)	148 (48.8%)	481 (52.8%)	898 (53.9%)
Monthly and less than monthly	80 (17.7%)	74 (24.4%)	209 (22.9%)	363 (21.8%)
Biweekly up to monthly	36 (8.0%)	24 (7.9%)	89 (9.8%)	149 (8.9%)
More than weekly up to biweekly	37 (8.2%)	36 (11.9%)	98 (10.8%)	171 (10.3%)
Unknown	29 (6.4%)	21 (6.9%)	34 (3.7%)	84 (5.0%)
# Medical care visits over 3 months	0.4 (1.3)	0.3 (1.0)	0.5 (1.4)	0.5 (1.3)
Medical care visit frequency (categorized)**				
None	345 (76.5%)	234 (77.2%)	667 (73.2%)	1246 (74.8%)
Monthly and less than monthly	56 (12.4%)	39 (12.9%)	174 (19.1%)	269 (16.2%)
Biweekly up to monthly	16 (3.5%)	9 (3.0%)	27 (3.0%)	52 (3.1%)
More than weekly up to biweekly	5 (1.1%)	0 (0.0%)	9 (1.0%)	14 (0.8%)
Unknown	29 (6.4%)	21 (6.9%)	34 (3.7%)	84 (5.0%)
Years of Pain***				
Less than 1 Year	83 (18.4%)	71 (7.8%)	56 (18.5%)	210 (12.6%)
1 Years to Less than 2 Years	36 (8.0%)	50 (5.5%)	18 (5.9%)	104 (6.2%)
2 Years to Less than 5 Years	80 (17.7%)	119 (13.1%)	50 (16.5%)	249 (15.0%)
5 Years to Less than 10 Years	71 (15.7%)	164 (18.0%)	50 (16.5%)	285 (17.1%)
10+ Years	176 (39.0%)	481 (52.8%)	119 (39.3%)	776 (46.6%)
Unknown	5 (1.1%)	26 (2.9%)	10 (3.3%)	41 (2.5%)
% of time spent in heavy labor				
None (0%)	220 (48.8%)	446 (49.0%)	158 (52.1%)	824 (49.5%)
Non-workplace: > 0% but < 20%	52 (11.5%)	108 (11.9%)	27 (8.9%)	187 (11.2%)
Workplace: > 0% but < 20%	90 (20.0%)	179 (19.6%)	55 (18.2%)	324 (19.5%)
Non-workplace: > 20%	14 (3.1%)	26 (2.9%)	7 (2.3%)	47 (2.8%)
Workplace: > 20%	45 (10.0%)	86 (9.4%)	29 (9.6%)	160 (9.6%)
Previous back surgery unsuccessful***	4 (0.9%)		1 (0.1%)	5 (0.4%)
Previous neck surgery unsuccessful		1 (0.3%)	3 (0.3%)	4 (0.3%)
Use of medications over past 6 months:				
Over-the-counter pain medications often/always	188 (41.7%)	392 (43.0%)	136 (44.9%)	716 (43.0%)

Table 1. Outcomes and patient characteristics predictors by type of chronic pain mean (SD) unless otherwise noted. (continued)

Label	CLBP (n = 451)	CLBP & CNP (n = 911)	CNP (n = 303)	Totals (n = 1665)
Prescription pain medications often/always	26 (5.8%)	77 (8.5%)	17 (5.6%)	120 (7.2%)
Narcotic medications often/always	21 (4.7%)	44 (4.8%)	8 (2.6%)	73 (4.4%)
Exercise in Past 6 Months: Often to Always*	313 (69.4%)	568 (62.3%)	210 (69.3%)	1091 (65.5%)
New patient (< 30 days)***	65 (14.4%)	67 (7.4%)	48 (15.8%)	180 (10.8%)
Ended chiropractic care during 3 mos***	55 (12.2%)	39 (4.3%)	17 (5.6%)	111 (6.7%)
Believe back pain is chronic	277 (61.4%)	569 (62.5%)		846 (62.1%)
Believe neck pain is chronic		575 (63.1%)	193 (63.7%)	768 (63.3%)
What low back pain would have been	6.4 (2.2)	6.5 (2.3)		6.5 (2.3)
What neck pain would have been		6.5 (2.4)	6.6 (2.3)	6.6 (2.4)
Exercise unsafe: Agree to strongly agree	24 (5.3%)	50 (5.5%)	12 (4.0%)	86 (5.2%)
Pain management self-efficacy (1-10)**	7.7 (1.8)	7.4 (1.8)	7.8 (1.7)	7.6 (1.8)
Expect chiropractic very-extremely successful	330 (73.2%)	658 (72.2%)	237 (78.2%)	1225 (73.6%)
Expect a lot to quite a bit of improvement	292 (64.7%)	569 (62.5%)	193 (63.7%)	1054 (63.3%)
Worry about pain: Mod to all the time**	72 (16.0%)	176 (19.3%)	36 (11.9%)	284 (17.1%)
Depression according to PROMIS items***	92 (20.4%)	260 (28.5%)	59 (19.5%)	411 (24.7%)
Affective distress (7-point scale)***	2.0 (1.1)	2.3 (1.2)	2.1 (1.2)	2.2 (1.2)
Catastrophizing (0-12 scale)***	2.0 (2.2)	2.5 (2.4)	1.8 (2.0)	2.3 (2.3)
Average age in years	48.4 (15.2)	46.4 (12.6)	47.4 (14.3)	47.5 (14.3)
Gender: Female***	265 (58.8%)	693 (76.1%)	245 (80.9%)	1203 (72.3%)
Education: Less than a 4-year degree*	194 (43.0%)	429 (47.1%)	118 (38.9%)	741 (44.5%)

CT = complementary therapy providers other than chiropractic—here, most often massage therapists; CLBP = chronic low back pain; CNP = chronic neck pain; LBP = low back pain; NP = neck pain

\*The values for the 3 pain categories are statistically different at  $P < 0.05$ .

\*\*The values for the 3 pain categories are statistically different at  $P < 0.01$ .

\*\*\*The values for the 3 pain categories are statistically different at  $P < 0.001$ .

had 6 chiropractic visits, 2 CT visits, and one-half medical provider visits. Less than half the sample had any CT visits (85% of these received massage, and about 25% each received physical therapy and/or acupuncture) and one-quarter had any medical provider visits (84% of these visited a general practitioner). Patients also consistently reported levels of what their pain would have been if they did not see their chiropractor, that were almost twice that of their current pain.

Tables 2 and 3 show the results of a series of HLM models that start as unconditional (time-only) models and then add chiropractic visits, other visits, and other characteristics as blocks of explanatory variables for each outcome. Tests of the unconditional models indicated that intercept (baseline) estimates varied significantly by patient and clinic, but weeks-since-baseline (time or slope) coefficient estimates varied only by patient and neither varied by region. The intercepts estimated for each unconditional model (Table 2) reflect the average baseline value for that outcome and the

estimates for the weeks-since-baseline coefficient show the average change in that outcome per week over the 3-month period. The estimated coefficients for weeks-since-baseline were all statistically significant and negative, indicating that on average these symptoms improve over time. Over the 3-month study period, ratings of low back pain were estimated to decrease an average of 0.03 points per week or 0.40 points over 3 months on a 0-10 scale. Ratings of neck pain decreased an average of 0.03 points per week or 0.44 points over 3 months. The ODI was estimated to decrease (improve) an average of 2.7 points (0.21\*13 weeks) and the NDI by 3.0 points (0.23\*13) over 3 months, both on a 0-100 scale.

The rows labeled clinic, ID, and residual partition the variance in the data for each model. The row labeled ID (intercept) had the largest value for each model, indicating that most of the variation in outcomes was due to variation in patients' baseline values. The small value given to ID (weeks) indicates that there

## Chiropractic Visit Frequency Impact on Outcomes

Table 2. Results of hierarchical linear models for each outcome: unconditional models and models of the impact of chiropractic visit frequency.

	LBP Rating (n = 1361)		ODI (n = 1362)		NP Rating (n = 1214)		NDI (n = 1213)	
	Unconditional	Adding visit frequency	Unconditional	Adding visit frequency	Unconditional	Adding visit frequency	Unconditional	Adding visit frequency
Intercept	3.48 (0.1)***	3.15 (0.1)***	19.27 (0.4)***	17.10 (0.6)***	3.64 (0.1)***	3.35 (0.1)***	21.37 (0.4)***	19.57 (0.7)***
Weeks Since Baseline (Weeks)	-0.03 (0.0)***	-0.02 (0.0)**	-0.21 (0.0)***	-0.22 (0.0)***	-0.03 (0.0)***	-0.02 (0.0)**	-0.23 (0.0)***	-0.18 (0.0)***
Visits to Chiropractor (Ref: Monthly to < monthly)								
Biweekly up to monthly	-	0.25 (0.1)	-	0.58 (0.8)	-	0.24 (0.1)	-	0.54 (0.9)
Weekly up to biweekly	-	0.60 (0.1)***	-	4.48 (0.9)***	-	0.48 (0.2)**	-	3.97 (0.9)***
More than weekly	-	0.95 (0.2)***	-	7.75 (1.1)***	-	0.72 (0.2)***	-	5.29 (1.2)***
Unknown	-	0.41 (0.3)	-	2.05 (1.6)	-	0.42 (0.3)	-	1.63 (1.8)
Weeks* Biweekly to monthly	-	0.00 (0.0)	-	0.08 (0.0)	-	-0.01 (0.0)	-	-0.07 (0.1)
Weeks*Weekly to biweekly	-	-0.01 (0.0)	-	0.05 (0.1)	-	-0.01 (0.0)	-	-0.07 (0.1)
Weeks*More than weekly	-	-0.05 (0.0)***	-	-0.14 (0.1)*	-	-0.03 (0.0)*	-	-0.14 (0.1)*
Partitioned Variance								
Clinic (Intercept)	0.04 (0.0)	0.04 (0.0)	5.26 (2.3)	5.79 (2.3)	0.09 (0.1)	0.10 (0.1)	5.23 (2.8)	5.77 (2.9)
ID (Weeks)	0.01 (0.0)	0.01 (0.0)	0.22 (0.0)	0.21 (0.0)	0.01 (0.0)	0.01 (0.0)	0.25 (0.0)	0.25 (0.0)
ID (Intercept)	2.51 (0.1)	2.40 (0.1)	122.00 (5.5)	114.81 (5.2)	2.82 (0.2)	2.75 (0.2)	127.04 (6.2)	122.85 (6.0)
Residual	1.61 (0.0)	1.61 (0.0)	27.75 (0.5)	27.75 (0.5)	1.60 (0.0)	1.60 (0.0)	32.67 (0.7)	32.67 (0.7)
D (df) for visits main effects	-	16.2 (4) ***	-	32.0 (4) ***	-	7.1 (4) **	-	14.7 (4) ***
D (df) for visits interactions	-	7.5 (3) **	-	6.3 (3) **	-	2.8 (3)	-	2.1 (3)

D = Deviance statistic with its degrees of freedom (number of parameters added over previous model), distributed  $\chi^2$ ; LBP = low back pain, 0-10 scale; NDI = Neck Disability Index, 0-100 scale; NP = neck pain, 0-10 scale; ODI = Oswestry Disability Index, 0-100 scale

\*The estimated coefficient or deviance statistic is statistically significant at  $P < 0.05$ .

\*\*The estimated coefficient or deviance statistic is statistically significant at  $P < 0.01$ .

\*\*\*The estimated coefficient or deviance statistic is statistically significant at  $P < 0.001$ .

was relatively little variance in the rate of improvement over time across patients. The clinic (intercept) terms indicate that there was some variance in patients' baseline values across clinics—i.e., clinics attract patients with different symptom severity. However, the weeks-since-baseline variance by clinic was not significant indicating that patients' improvement over time did not vary by clinic. The residual indicates the amount of unexplainable variance.

The significance of the coefficients estimated when adding chiropractic visit frequency to the unconditional models in Table 2, indicate that having more frequent visits is associated with higher levels of pain and disability at baseline (main effects), but only those who see their chiropractor more than weekly

had significantly faster improvement (interactions with weeks). The deviance statistics shown in the last 2 rows indicate the significance of the explanatory power of adding each block of variables to models containing all previous variables. Adding chiropractic visit frequency main effects to the unconditional models (i.e., allowing chiropractic visit frequency to explain baseline symptoms) provided a significant amount of explanatory power to all models. However, adding chiropractic visit frequency interactions with time (weeks) only provided significant explanatory power (i.e., was associated with more improvement in outcomes) for CLBP, but not CNP alone.

Table 3 shows the estimated coefficients and deviance statistics for adding main effects and interaction

Table 3. Results of hierarchical linear models for each outcome: models adding the impact of frequency of other types of visits and other patient characteristics.

	LBP Rating (n = 1361)		ODI (n = 1362)		NP Rating (n = 1214)		NDI (n = 1213)	
	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables
Intercept	2.82 (0.1)***	0.91 (0.3)**	14.81 (0.6)***	11.80 (2.1)***	3.05 (0.1)***	0.10 (0.4)	17.41 (0.7)***	4.16 (2.3)
Weeks Since BL (Weeks)	-0.02 (0.0)**	-0.02 (0.0)**	-0.25 (0.0)***	-0.25 (0.0)***	-0.02 (0.0)*	-0.02 (0.0)*	-0.21 (0.0)***	-0.21 (0.0)***
Visits to chiropractor (Ref: Monthly to < monthly)								
Biweekly up to monthly	0.24 (0.1)	0.19 (0.1)	0.66 (0.8)	0.51 (0.6)	0.25 (0.1)	0.21 (0.1)*	0.68 (0.9)	0.69 (0.7)
Weekly up to biweekly	0.55 (0.1)***	0.22 (0.1)*	4.13 (0.8)***	2.32 (0.6)***	0.45 (0.1)**	0.19 (0.1)	3.83 (0.9)***	2.53 (0.7)***
More than weekly	0.82 (0.2)***	0.40 (0.1)**	6.59 (1.0)***	3.95 (0.8)***	0.62 (0.2)***	0.24 (0.1)	4.14 (1.1)***	2.18 (0.9)*
Unknown	0.74 (0.3)**	0.51 (0.2)*	4.33 (1.6)**	2.06 (1.3)	0.70 (0.3)*	0.58 (0.2)*	3.69 (1.8)*	3.61 (1.4)*
Wks*Biweekly to monthly	0.00 (0.0)	0.00 (0.0)	0.08 (0.0)	0.08 (0.0)	-0.01 (0.0)	-0.01 (0.0)	-0.06 (0.1)	-0.06 (0.1)
Wks*Weekly to biweekly	-0.01 (0.0)	-0.01 (0.0)	0.04 (0.1)	0.04 (0.1)	-0.01 (0.0)	-0.01 (0.0)	-0.06 (0.1)	-0.06 (0.1)
Wks*More than weekly	-0.05 (0.0)***	-0.05 (0.0)***	-0.16 (0.1)*	-0.16 (0.1)*	-0.03 (0.0)*	-0.03 (0.0)*	-0.16 (0.1)*	-0.16 (0.1)*
Other CT provider visits								
Monthly to < monthly	0.37 (0.1)**	0.22 (0.1)*	0.61 (0.8)	0.16 (0.6)	0.40 (0.1)**	0.23 (0.1)*	1.46 (0.8)	0.55 (0.7)
Biweekly up to monthly	0.42 (0.2)*	0.19 (0.1)	2.21 (1.1)*	1.43 (0.8)	0.33 (0.2)	0.10 (0.1)	0.21 (1.2)	-1.23 (0.9)
> Weekly up to biweekly	0.64 (0.2)***	0.33 (0.1)*	5.66 (1.0)***	4.43 (0.8)***	0.70 (0.2)***	0.45 (0.1)**	4.31 (1.1)***	2.89 (0.9)***
Unknown	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)
Wks* Monthly to <monthly	0.00 (0.0)	0.00 (0.0)	0.04 (0.0)	0.04 (0.0)	-0.01 (0.0)	-0.01 (0.0)	-0.11 (0.1)*	-0.11 (0.1)*
Wks* Biweekly to monthly	-0.01 (0.0)	-0.01 (0.0)	0.00 (0.1)	0.00 (0.1)	0.01 (0.0)	0.01 (0.0)	0.13 (0.1)	0.14 (0.1)
Wks*>Weekly to biweekly	0.01 (0.0)	0.01 (0.0)	0.12 (0.1)	0.12 (0.1)	0.00 (0.0)	0.00 (0.0)	0.11 (0.1)	0.11 (0.1)
Medical provider visit								
Monthly to < monthly	0.58 (0.1)***	0.20 (0.1)	4.75 (0.8)***	1.78 (0.6)**	0.35 (0.1)*	-0.06 (0.1)	4.20 (0.9)***	1.17 (0.7)
Biweekly up to monthly	1.46 (0.3)***	0.78 (0.2)***	15.40 (1.7)***	8.36 (1.3)***	1.41 (0.3)***	0.73 (0.3)**	17.24 (2.0)***	10.02 (1.6)***
> Weekly up to biweekly	1.09 (0.5)*	0.35 (0.4)	13.10 (2.9)***	5.92 (2.3)**	0.30 (0.6)	-0.18 (0.5)	6.79 (3.9)	0.00 (3.1)
Unknown	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)
Wks* Monthly to <monthly	0.00 (0.0)	0.00 (0.0)	0.01 (0.1)	0.01 (0.1)	0.00 (0.0)	0.00 (0.0)	0.12 (0.1)*	0.12 (0.1)*
Wks* Biweekly to monthly	-0.02 (0.0)	-0.02 (0.0)	0.17 (0.1)	0.17 (0.1)	-0.01 (0.0)	-0.01 (0.0)	-0.08 (0.1)	-0.08 (0.1)
Wks*>Weekly to biweekly	-0.01 (0.0)	-0.02 (0.0)	0.07 (0.2)	0.06 (0.2)	0.10 (0.1)	0.10 (0.1)	0.99 (0.3)***	0.99 (0.3)***
Other patient characteristics	See appendix for coefficients		See appendix for coefficients		See appendix for coefficients		See appendix for coefficients	
Partitioned Variance								
Clinic (Intercept)	0.02 (0.0)	0.01 (0.0)	5.48 (2.1)	1.84 (1.1)	0.07 (0.0)	0.00 (0.0)	3.80 (2.3)	0.57 (1.3)
ID (Weeks)	0.01 (0.0)	0.01 (0.0)	0.21 (0.0)	0.21 (0.0)	0.01 (0.0)	0.01 (0.0)	0.23 (0.0)	0.24 (0.0)



## Chiropractic Visit Frequency Impact on Outcomes

Table 3. Results of hierarchical linear models for each outcome: models adding the impact of frequency of other types of visits and other patient characteristics. (continued)

	LBP Rating (n = 1361)		ODI (n = 1362)		NP Rating (n = 1214)		NDI (n = 1213)	
	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables
ID (Intercept)	2.23 (0.1)	1.02 (0.1)	99.93 (4.6)	52.38 (2.7)	2.64 (0.1)	1.13 (0.1)	111.07 (5.5)	58.86 (3.4)
Residual	1.61 (0.0)	1.61 (0.0)	27.76 (0.5)	27.73 (0.5)	1.60 (0.0)	1.60 (0.0)	32.65 (0.7)	32.62 (0.7)
D (df) for main effects	45.9 (8) ***	300.7 (33) ***	92.1 (8) ***	402.0 (33) ***	25.6 (8) ***	288.4 (33) ***	64.5 (8) ***	355.9 (33) ***
D (df) for interactions	0.8 (6)	-61.5 (32)	3.6 (6)	-25.4 (32)	2.5 (6)	-54.4 (32)	15.3 (6) ***	-12.2 (32)

BL = Baseline; CT = complementary therapy providers—here, most often massage therapists; D = Deviance statistic with its degrees of freedom (number of parameters added over previous model), distributed  $\chi^2$ ; LBP = low back pain, 0-10 scale; NDI = Neck Disability Index, 0-100 scale; NP = neck pain, 0-10 scale; ODI = Oswestry Disability Index, 0-100 scale; OTC = Over-the-counter medications; Rx = Prescription medications

\*The estimated coefficient or deviance statistic is statistically significant at  $p < .05$ .

\*\*The estimated coefficient or deviance statistic is statistically significant at  $p < .01$ .

\*\*\*The estimated coefficient or deviance statistic is statistically significant at  $p < .001$ .

terms for other types of visits and then for all other explanatory variables. The estimated coefficients for the other explanatory (nonvisit) variables are shown in the Appendix Table 1. Because the deviance statistics indicated that model fit was not improved by adding interaction terms for all other explanatory variables, coefficients reported for these variables in the Appendix Table 1 are from models without these interactions. Note that the size of the main effect coefficients for chiropractic visit frequency diminish somewhat after adding all explanatory variables, but the size and significance of the interaction coefficients remain fairly constant.

The coefficients and deviance statistics for adding the effects of other types of visits indicate that they were associated with all baseline outcomes, but only associated with changes in neck function (NDI) over time. Similar to chiropractic visits, the main effects coefficients were positive and generally increased across frequency categories, indicating that more visits were associated with higher (worse) baseline outcomes. Using CT visits monthly to less than monthly was associated with increased improvement in neck function. However, in contrast, the positive significant coefficients for interaction terms for medical provider visits for neck function indicate that patients with those levels of medical visits had less improvement than was seen with patients who did not see medical providers. Adding in all explanatory variables reduced the size of the main effects but had no effect on the interactions.

The partitioned variance statistics at the bottom of Table 3 indicate that these models were able to explain

almost all the variance seen in patients' baseline values by clinic and over half the nonclinic-based baseline variance. However, these models reduced little of the variance seen in patients' improvement over time.

### Discussion

Our results raise interesting considerations for coverage policies for chronic spinal pain, including visit frequencies associated with better outcomes and appropriate care after patient improvement has plateaued (reached MTI).

If the main goal of patients and clinicians is better symptom improvement, rather than maintaining current symptoms, these data indicate that this might require more than once-per-week chiropractic visits for those with CLBP and possibly the addition of massage to chiropractic care for CNP functional improvement. The more-than-weekly chiropractic visit frequency associated with increased improvement occurred more often in patients with worse baseline pain and function who may have had more room for improvement. Nevertheless, in this sample, further symptom improvement may not be the main goal (39).

The slight improvement in symptoms over time and the small variance in that improvement across patients may indicate that most of these patients' symptoms have plateaued at (or near) their MTI. Once MTI is reached, treatment focus changes from symptom improvement to maintenance and/or management. Therefore, policies that require documentation of ongoing clinical improvement for continued care may not be appropriate (21,23-26). This finding of patients

reaching MTI is consistent with what other studies have shown for this sample—that they value maintenance of their present symptom levels and that most have a goal of pain management, not for a cure (39,40).

While the majority saw their chiropractor every 2 weeks at most, patients managed their pain using a variety of different visit frequencies. In previous analyses of these data, we found that chiropractic visit frequency was predicted by patients' baseline function, stage of care (whether a new patient or near ending care), and the characteristics of the treating chiropractor (36).

Finally, while most guidelines agree that continued treatment (e.g., chronic pain management or support care) may be needed under certain conditions after MTI is reached, little information is available to determine the treatment appropriate to maintain symptom gains (21,22). Some guidelines have suggested that documentation of clinical deterioration with treatment withdrawal be required to identify those who need ongoing care (21,25,26). Five points are offered regarding appropriate ongoing care. First, given that these patients have had their pain an average of 14 years and have used chiropractic care for 11 years, consideration must be given to the burden of repeated treatment withdrawals and their requirement to qualify for continued care (29). Second, only 6.7% of this sample ended care (and half of these also restarted care) during our study period [data not shown]. These may not be formal treatment withdrawals, but 70% of these patients reported they ended care because they were better and no longer needed treatment (most others ended because of lost insurance coverage or relocation). Third, patients reported (0-10 scale) what they believed their pain would be if they did not see their chiropractor and these reports were about 3 points above current pain—more than the 2-point minimal clinically important change for pain suggested by one guideline to justify ongoing care (21,50). Although these reports could be based on psychosocial factors such as fear/anxiety regarding not receiving treatment, given the length of time these patients have had their pain, they could also be based on lived experience with past treatment withdrawals. Fourth, because these patients reported at baseline that their current symptoms were mild (average pain intensity of 3 to 4 on a 0-10 scale with minimal-to-moderate back dysfunction and mild neck dysfunction), but still improving over time, it could be argued that they were successfully managing their

CLBP and CNP (29,46,55). Fifth, given this successful management using a variety of visit frequencies, it could be argued that each individual be covered as needed for visits. This need can ebb and flow, and is tempered by patients' out-of-pocket cost of care: even with some insurance coverage, a visit to a chiropractor (or any recommended nonpharmacologic therapy provider) is usually associated with a per-visit out-of-pocket co-payment in addition to the cost of travel to the visit and of missing work (83-85).

This study benefits from a large longitudinal sample of chronic pain patients, but it also has limitations. Our sample may not be representative of all patients with CLBP and CNP, but it is representative of chiropractic patients with CLBP and CNP in terms of age, gender, race/ethnicity, income, education, and insurance coverage for chiropractic (28,29,31,86-88). It is also representative of pain and function levels seen in patients under treatment for CLBP and CNP (37,38). Although similar demographic profiles have also been found for those using other nonpharmacologic therapies for spinal problems, our study's results should not be generalized to patients who are not using these therapies now (28). Our data were self-report and may be subject to response (e.g., social desirability, recall) biases. Our study was observational; although associations between visits and other key variables, and outcomes and their improvement have been shown, without randomization and a control group we cannot say whether a change in allowed visit frequency would make a difference in these patients' choices and outcomes. Our sample excluded patients with work-related injuries or personal injury claims. We did not capture specific treatments received during visits, which could affect outcomes. Finally, our data were restricted to a 3-month window into symptoms and care for a chronic condition. Even though both were fairly consistent over these months, a longer period may have shown different patterns.

It seems that some long-term CLBP and CNP patients may be successfully managing (and slightly improving) their chronic pain while using chiropractic care. These patients do this using a variety of visit frequencies. Treatment algorithms requiring demonstration of continued clinical improvement seem inconsistent with successful pain management, especially if patients have reached a plateau, and requirements of repeated demonstrations of symptom deterioration with treatment withdrawal seem unethical, especially for those with long-term chronic pain. Nevertheless, payers clearly need evidence to support new coverage policies for on-

going nonpharmacologic care for patients with chronic pain, including chiropractic care (85,89). This study may illustrate an example of successful nonpharmacologic pain management that deserves further consideration from a policy perspective. In addition, future studies are needed to clarify the impact of various chiropractic coverage policies on clinical outcomes and costs.

### Author Contributions

Drs. Coulter and Herman and Ms. Edgington had

full access to all of the data in the study and jointly take responsibility for the integrity of the data. Dr. Herman and Ms. Edgington take responsibility for the accuracy of the data analysis. Drs. Herman, Goertz, Hurwitz, and Sorbero designed the study protocol. Dr. Herman managed the literature search and summary of previous related work and wrote the first draft of the manuscript. Drs. Coulter, Edgington, Goertz, Hurwitz and Sorbero provided revision for intellectual content and final approval of the manuscript.

### REFERENCES

- Institute of Medicine. Relieving pain in America: A blueprint for transforming prevention, care, education, and research. The National Academies Press, Washington, DC, 2011.
- Johannes CB, Le TK, Zhou X, Johnston JA, Dworkin RH. The prevalence of chronic pain in United States adults: Results of an internet-based survey. *J Pain* 2010; 11:1230-1239.
- Martin BI, Deyo RA, Mirza SK, et al. Expenditures and health status among adults with back and neck problems. *JAMA* 2008; 299:656-664.
- Martin BI, Turner JA, Mirza SK, Lee MJ, Comstock BA, Deyo RA. Trends in health care expenditures, utilization, and health status among US adults with spine problems, 1997-2006. *Spine* 2009; 34:2077-2084.
- Gore M, Sadosky A, Stacey BR, Tai K-S, Leslie D. The burden of chronic low back pain: Clinical comorbidities, treatment patterns, and health care costs in usual care settings. *Spine* 2012; 37:E668-E677.
- Geurts JW, Willems PC, Kallewaard J-W, van Kleef M, Dirksen C. The impact of chronic discogenic low back pain: Costs and patients' burden. *Pain Res Manage* 2018; 2018:1-8.
- Compare A, Marchettini P, Zarbo C. Risk factors linked to psychological distress, productivity losses, and sick leave in low-back-pain employees: A three-year longitudinal cohort study. *Pain Res Treatment* 2016; 2016:1-9.
- Smith M, Davis MA, Stano M, Whedon JM. Aging baby boomers and the rising cost of chronic back pain: Secular trend analysis of longitudinal Medical Expenditures Panel Survey data for years 2000 to 2007. *J Manipulative Physiol Ther* 2013; 36:2-11.
- Dieleman JL, Baral R, Birger M, et al. US spending on personal health care and public health, 1996-2013. *JAMA* 2016; 316:2627-2646.
- Stewart WF, Ricci JA, Chee E, Morganstein D, Lipton R. Lost productive time and cost due to common pain conditions in the US workforce. *JAMA* 2003; 290:2443-2454.
- Ivanova JI, Birnbaum HG, Schiller M, Kantor E, Johnstone BM, Swindle RW. Real-world practice patterns, health-care utilization, and costs in patients with low back pain: The long road to guideline-concordant care. *Spine J* 2011; 11:622-632.
- Gore M, Tai KS, Sadosky A, Leslie D, Stacey BR. Use and costs of prescription medications and alternative treatments in patients with osteoarthritis and chronic low back pain in community-based settings. *Pain Pract* 2012; 12:550-560.
- Qaseem A, Wilt TJ, McLean RM, Forcica MA. Noninvasive treatments for acute, subacute, and chronic low back pain: A clinical practice guideline from the American College of Physicians. *Ann Intern Med* 2017; 166:514-530.
- The Diagnosis and Treatment of Low Back Pain Work Group. VA/DoD Clinical practice guideline for diagnosis and treatment of low back pain, version 2.0. The Office of Quality, Safety and Value, VA, & Office of Evidence Based Practice, U.S. Army Medical Command, Washington, DC, 2017. [www.healthquality.va.gov/guidelines/Pain/lbp/](http://www.healthquality.va.gov/guidelines/Pain/lbp/). Accessed 3/28/2018.
- The Joint Commission. Non-pharmacologic and non-opioid solutions for pain management. *Quick Safety* 2018; August:1-2. [www.jointcommission.org/resources/news-and-multimedia/newsletters/newsletters/quick-safety/quick-safety-44-nonpharmacologic-and-nonopioid-solutions-for-pain-management/](http://www.jointcommission.org/resources/news-and-multimedia/newsletters/newsletters/quick-safety/quick-safety-44-nonpharmacologic-and-nonopioid-solutions-for-pain-management/).
- U.S. Department of Health and Human Services. Pain Management Best Practices Inter-Agency Task Force report: Updates, gaps, inconsistencies, and recommendations. U. S. Department of Health and Human Services, Washington, DC, 2019. [www.hhs.gov/ash/advisory-committees/pain/reports/index.html](http://www.hhs.gov/ash/advisory-committees/pain/reports/index.html). Accessed 6/19/2019.
- Chronic Pain: Symptoms, Diagnosis, & Treatment, *NIH Medline Plus*. National Institutes of Health and Friends of the National Library of Medicine, Bethesda, MD, 2011, pp 5-6.
- Ballantyne JC, Shin NS. Efficacy of opioids for chronic pain: A review of the evidence. *Clin J Pain* 2008; 24:469-478.
- Ling W, Mooney L, Hillhouse M. Prescription opioid abuse, pain and addiction: Clinical issues and implications. *Drug Alcohol Rev* 2011; 30:300-305.
- California Division of Workers' Compensation. Medical Treatment Utilization Schedule (MTUS) Chronic Pain Medical Treatment Guidelines State of California, Sacramento,

- CA, 2016. [www.dir.ca.gov/dwc/DWCPropRegs/MTUS-Opioids-ChronicPain/Final-Regulations/CleanCopy/Chronic-Pain-Guidelines.pdf](http://www.dir.ca.gov/dwc/DWCPropRegs/MTUS-Opioids-ChronicPain/Final-Regulations/CleanCopy/Chronic-Pain-Guidelines.pdf). Accessed 07/12/2019.
21. Farabaugh RJ, Dehen MD, Hawk C. Management of chronic spine-related conditions: Consensus recommendations of a multidisciplinary panel. *J Manipulative Physiol Ther* 2010; 33:484-492.
  22. Canadian Chiropractic Association; Canadian Federation of Chiropractic Regulatory Boards; Clinical Practice Guidelines Development Initiative; et al. Chiropractic clinical practice guideline: Evidence-based treatment of adult neck pain not due to whiplash. *J Canadian Chiropractic Assoc* 2005; 49:158-209.
  23. Colorado Division of Workers' Compensation. Low back pain medical treatment guidelines. Colorado Division of Workers' Compensation, Denver, CO, 2014, pp 112. [https://cdle.colorado.gov/sites/cdle/files/MTG\\_Ex1\\_LBP.pdf](https://cdle.colorado.gov/sites/cdle/files/MTG_Ex1_LBP.pdf). Accessed 02/01/2018.
  24. Colorado Division of Workers' Compensation. Cervical spine injury medical treatment guidelines. Colorado Division of Workers' Compensation, Denver, CO, 2014, pp 96. [https://cdle.colorado.gov/sites/cdle/files/MTG\\_Ex8\\_CSI.pdf](https://cdle.colorado.gov/sites/cdle/files/MTG_Ex8_CSI.pdf). Accessed 02/01/2018.
  25. Globe GA, Morris CE, Whalen WM, Farabaugh RJ, Hawk C. Chiropractic management of low back disorders: Report from a consensus process. *J Manipulative Physiol Ther* 2008; 31:651-658.
  26. Globe G, Farabaugh RJ, Hawk C, et al. Clinical practice guideline: Chiropractic care for low back pain. *J Manipulative Physiol Ther* 2016; 39:1-22.
  27. Hurwitz EL. Epidemiology: Spinal manipulation utilization. *J Electromyography Kinesiol* 2012; 22:648-654.
  28. Martin BI, Gerkovich MM, Deyo RA, et al. The association of complementary and alternative medicine use and health care expenditures for back and neck problems. *Med Care* 2012; 50:1029-1036.
  29. Herman PM, Kommareddi M, Sorbero ME, et al. Characteristics of chiropractic patients being treated for chronic low back and chronic neck pain. *J Manipulative Physiol Ther* 2018; 41:445-455.
  30. Cooper RA, McKee HJ. Chiropractic in the United States: Trends and issues. *Milbank Q* 2003; 81:107-138.
  31. Coulter ID, Shekelle PG. Chiropractic in North America: A descriptive analysis. *J Manipulative Physiol Ther* 2005; 28:83-89.
  32. Posner J, Glew C. Neck pain. *Ann Intern Med* 2002; 136:758-759.
  33. Gemmell HA, Hayes BM. Patient satisfaction with chiropractic physicians in an independent physicians' association. *J Manipulative Physiol Ther* 2001; 24:556-559.
  34. MacPherson H, Newbronner E, Chamberlain R, Hopton A. Patients' experiences and expectations of chiropractic care: A national cross-sectional survey. *Chiropr Man Therap* 2015; 23:3.
  35. Penney LS, Ritenbaugh C, DeBar LL, Elder C, Deyo RA. Provider and patient perspectives on opioids and alternative treatments for managing chronic pain: A qualitative study. *BMC Fam Pract* 2017; 17:164.
  36. Herman PM, Edgington SE, Hurwitz EL, Coulter ID. Predictors of visit frequency for patients using ongoing chiropractic care for chronic low back and chronic neck pain. *BMC Musculoskel Disord* 2020; 21:298.
  37. Evans R, Bronfort G, Nelson B, Goldsmith CH. Two-year follow-up of a randomized clinical trial of spinal manipulation and two types of exercise for patients with chronic neck pain. *Spine* 2002; 27:2383-2389.
  38. Niemistö L, Lahtinen-Suopanki T, Rissanen P, Lindgren K-A, Sarna S, Hurri H. A randomized trial of combined manipulation, stabilizing exercises, and physician consultation compared to physician consultation alone for chronic low back pain. *Spine* 2003; 28:2185-2191.
  39. Herman PM, Edgington SE, Ryan GW, Coulter ID. Prevalence and characteristics of chronic spinal pain patients with different hopes (treatment goals) for ongoing chiropractic care. *J Altern Complement Med* 2019; 25:1015-1025.
  40. Herman PM, Luoto JE, Kommareddi M, Sorbero ME, Coulter ID. Patient willingness to pay (WTP) for reductions in chronic low back pain and chronic neck pain. *J Pain* 2019; 20:1317-1327.
  41. Coulter ID, Herman PM, Ryan GW, Hays RD, Hilton LG, Whitley MD. Researching the appropriateness of care in the complementary and integrative health professions: Part 1. *J Manipulative Physiol Ther* 2018; 41:800-806.
  42. Coulter ID, Herman PM, Ryan GW, Hays RD, Hilton LJ, CERC Team. The challenge of determining appropriate care in the era of patient-centered care and rising health care costs. *J Health Serv Res Policy* 2019; 24:201-206.
  43. Coulter ID, Aliyev GR, Whitley MD, et al. Researching the appropriateness of care in the complementary and integrative health professions Part 4: Putting practice back into evidence-based practice by recruiting clinics and patients. *J Manipulative Physiol Ther* 2019; 42:319-326.
  44. Huskisson E. Measurement of pain. *Lancet* 1974; 2:1127-1131.
  45. Vernon H, Mior S. The Neck Disability Index: A study of reliability and validity. *J Manipulative Physiol Ther* 1991; 14:409-415.
  46. Fairbank JCT, Couper J, Davies JB, O'Brien JP. The Oswestry low back pain disability questionnaire. *Physiotherapy* 1980; 66:271-273.
  47. Bijur PE, Latimer CT, Gallagher EJ. Validation of a verbally administered numerical rating scale of acute pain for use in the emergency department. *Acad Emerg Med* 2003; 10:390-392.
  48. Childs JD, Piva SR, Fritz JM. Responsiveness of the numeric pain rating scale in patients with low back pain. *Spine* 2005; 30:1331-1334.
  49. Paice JA, Cohen FL. Validity of a verbally administered numeric rating scale to measure cancer pain intensity. *Cancer Nurs* 1997; 20:88-93.
  50. Salaffi F, Stancati A, Silvestri CA, Ciapetti A, Grassi W. Minimal clinically important changes in chronic musculoskeletal pain intensity measured on a numerical rating scale. *Eur J Pain* 2004; 8:283-291.
  51. Downie W, Leatham P, Rhind V, Wright V, Branco J, Anderson J. Studies with pain rating scales. *Ann Rheum Dis* 1978; 37:378-381.
  52. Cleland JA, Fritz JM, Whitman JM, Palmer JA. The reliability and construct validity of the Neck Disability Index and patient specific functional scale in patients with cervical radiculopathy. *Spine* 2006; 31:598-602.
  53. Wheeler AH, Goolkasian P, Baird AC, Darden BV. Development of the Neck Pain and Disability Scale: Item analysis, face, and criterion-related validity. *Spine* 1999; 24:1290-1294.
  54. McCarthy M, Grevitt M, Silcocks P,

- Hobbs G. The reliability of the Vernon and Mior neck disability index, and its validity compared with the short form-36 health survey questionnaire. *Eur Spine J* 2007; 16:2111-2117.
55. Vernon H. The Neck Disability Index: State-of-the-art, 1991-2008. *J Manipulative Physiol Ther* 2008; 31:491-502.
56. Davidson M, Keating JL. A comparison of five low back disability questionnaires: Reliability and responsiveness. *Phys Ther* 2002; 82:8-24.
57. Fisher K, Johnston M. Validation of the Oswestry low back pain disability questionnaire: Its sensitivity as a measure of change following treatment and its relationship with other aspects of the chronic pain experience. *Physiother Theory Pract* 1997; 13:67-80.
58. Grönblad M, Hupli M, Wennerstrand P, et al. Inter-correlation and test-retest reliability of the pain disability index (PDI) and the Oswestry disability questionnaire (ODQ) and their correlation with pain intensity in low back pain patients. *Clin J Pain* 1993; 9:189-195.
59. Schellingerhout JM, Verhagen AP, Heymans MW, et al. Which subgroups of patients with non-specific neck pain are more likely to benefit from spinal manipulation therapy, physiotherapy, or usual care? *Pain* 2008; 139:670-680.
60. Underwood M, Morton V, Farrin A, Team UBT. Do baseline characteristics predict response to treatment for low back pain? Secondary analysis of the UK BEAM dataset [ISRCTN32683578]. *Rheumatology* 2007; 46:1297-1302.
61. Cook CE, Learman KE, O'halloran BJ, et al. Which prognostic factors for low back pain are generic predictors of outcome across a range of recovery domains? *Phys Ther* 2013; 93:32-40.
62. Jensen OK, Nielsen CV, Stengaard-Pedersen K. One-year prognosis in sick-listed low back pain patients with and without radiculopathy. Prognostic factors influencing pain and disability. *Spine J* 2010; 10:659-675.
63. Rasmussen-Barr E, Bohman T, Hallqvist J, Holm L, Skillgate E. Do physical activity level and body mass index predict recovery from persistent neck pain in men and women of working age? A population-based cohort study. *Eur Spine J* 2013; 22:2077-2083.
64. Riipinen M, Niemisto L, Lindgren K-A, Hurri H. Psychosocial differences as predictors for recovery from chronic low back pain following manipulation, stabilizing exercises and physician consultation or physician consultation alone. *J Rehabil Med* 2005; 37:152-158.
65. Moffett JAK, Carr J, Howarth E. High fear-avoiders of physical activity benefit from an exercise program for patients with back pain. *Spine* 2004; 29:1167-1172.
66. Verkerk K, Luijsterburg P, Heymans M, et al. Prognosis and course of pain in patients with chronic non-specific low back pain: A 1-year follow-up cohort study. *Eur J Pain* 2015; 19:1101-1110.
67. Smeets R, Beelen S, Goossens M, Schouten E, Knottnerus J, Vlaeyen JWS. Treatment expectancy and credibility are associated with the outcome of both physical and cognitive-behavioral treatment in chronic low back pain: In Smeets R (ed). Active rehabilitation for chronic low back pain: Cognitive-behavioral, physical, or both? Maastricht University, Eindhoven, The Netherlands 2008, pp 138-160.
68. Smeets RJ, Maher CG, Nicholas MK, Refshauge KM, Herbert RD. Do psychological characteristics predict response to exercise and advice for subacute low back pain? *Arthritis Rheum* 2009; 61:1202-1209.
69. Niemisto L, Sarna S, Lahtinen-Suopanki T, Lindgren K-A, Hurri H. Predictive factors for 1-year outcome of chronic low back pain following manipulation, stabilizing exercises, and physician consultation for physician consultation alone. *J Rehabil Med* 2004; 36:104-109.
70. Hill JC, Lewis M, Sim J, Hay EM, Dziedzic K. Predictors of poor outcome in patients with neck pain treated by physical therapy. *Clin J Pain* 2007; 23:683-690.
71. Smeets RJ, Vlaeyen JW, Kester AD, Knottnerus JA. Reduction of pain catastrophizing mediates the outcome of both physical and cognitive-behavioral treatment in chronic low back pain. *J Pain* 2006; 7:261-271.
72. van der Windt DA, Kuijpers T, Jellema P, van der Heijden GJ, Bouter LM. Do psychological factors predict outcome in both low-back pain and shoulder pain? *Ann Rheum Dis* 2007; 66:313-319.
73. Anderson KO, Dowds BN, Pelletz RE, Edwards WT, Peeters-Asdourian C. Development and initial validation of a scale to measure self-efficacy beliefs in patients with chronic pain. *Pain* 1995; 63:77-84.
74. Devilly GJ, Borkovec TD. Psychometric properties of the credibility/expectancy questionnaire. *J Behav Ther Exp Psychiatry* 2000; 31:73-86.
75. Amtmann D, Kim J, Chung H, et al. Comparing CESD-10, PHQ-9, and PROMIS depression instruments in individuals with multiple sclerosis. *Rehabil Psychol* 2014; 59:220-229.
76. Kerns RD, Turk DC, Rudy TE. The West Haven-Yale multidimensional pain inventory (WHYMPI). *Pain* 1985; 23:345-356.
77. Sullivan MJ, Bishop SR, Pivik J. The pain catastrophizing scale: Development and validation. *Psychol Assess* 1995; 7:524-532.
78. Heo M, Faith MS, Mott JW, Gorman BS, Redden DT, Allison DB. Hierarchical linear models for the development of growth curves: An example with body mass index in overweight/obese adults. *Stat Med* 2003; 22:1911-1942.
79. Raudenbush SW, Bryk AS. *Hierarchical linear models: Applications and data analysis methods second edition*. Sage Publications, Thousand Oaks, CA, 2002.
80. Singer JD. Using SAS PROC MIXED to fit multilevel models, hierarchical models, and individual growth models. *J Educ Behav Stat* 1998; 23:323-355.
81. Feldman HA. Families of lines: Random effects in linear regression analysis. *J Appl Physiol* 1988; 64:1721-1732.
82. O'Connell AA, McCoach DB. Applications of hierarchical linear models for evaluations of health interventions: Demystifying the methods and interpretations of multilevel models. *Eval Health Prof* 2004; 27:119-151.
83. McGorry RW, BSPT BSW, Snook SH, Hsiang SM. The relation between pain intensity, disability, and the episodic nature of chronic and recurrent low back pain. *Spine* 2000; 25:834-841.
84. Cassidy JD, Cote P, Carroll LJ, Kristman V. Incidence and course of low back pain episodes in the general population. *Spine* 2005; 30:2817-2823.
85. Heyward J, Jones CM, Compton WM, et al. Coverage of nonpharmacologic treatments for low back pain among US public and private insurers. *JAMA Netw Open* 2018; 1:e183044.
86. Mootz RD, Cherkin DC, Odegard CE, Eisenberg DM, Barassi JP, Deyo RA. Characteristics of chiropractic practitioners, patients, and encounters

- in Massachusetts and Arizona. *J Manipulative Physiol Ther* 2005; 28:645-653.
87. Stevans JM, Zodet MW. Clinical, demographic, and geographic determinants of variation in chiropractic episodes of care for adults using the 2005-2008 Medical Expenditure Panel Survey. *J Manipulative Physiol Ther* 2012; 35:589-599.
88. Hurwitz EL, Coulter ID, Adams AH, Genovese BJ, Shekelle PG. Use of chiropractic services from 1985 through 1991 in the United States and Canada. *Am J Public Health* 1998; 88:771-776.
89. Goertz CM, George SZ. Insurer coverage of nonpharmacological treatments for low back pain—Time for a change. *JAMA Netw Open* 2018; 1:e183037.

## Chiropractic Visit Frequency Impact on Outcomes

Appendix Table 1. Results of hierarchical linear models for each outcome: models adding the impact of frequency of other types of visits and other patient characteristics now showing the estimated coefficients for other patient characteristics not included in Table 3.

	LBP Rating (n = 1361)		ODI (n = 1362)		NP Rating (n = 1214)		NDI (n = 1213)	
	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables
Intercept	2.82 (0.1)***	0.91 (0.3)**	14.81 (0.6)***	11.80 (2.1)***	3.05 (0.1)***	0.10 (0.4)	17.41 (0.7)***	4.16 (2.3)
Weeks Since BL (Weeks)	-0.02 (0.0)**	-0.02 (0.0)**	-0.25 (0.0)***	-0.25 (0.0)***	-0.02 (0.0)*	-0.02 (0.0)*	-0.21 (0.0)***	-0.21 (0.0)***
Visits to chiropractor (Ref: Monthly to < monthly)								
Biweekly up to monthly	0.24 (0.1)	0.19 (0.1)	0.66 (0.8)	0.51 (0.6)	0.25 (0.1)	0.21 (0.1)*	0.68 (0.9)	0.69 (0.7)
Weekly up to biweekly	0.55 (0.1)***	0.22 (0.1)*	4.13 (0.8)***	2.32 (0.6)***	0.45 (0.1)**	0.19 (0.1)	3.83 (0.9)***	2.53 (0.7)***
More than weekly	0.82 (0.2)***	0.40 (0.1)**	6.59 (1.0)***	3.95 (0.8)***	0.62 (0.2)***	0.24 (0.1)	4.14 (1.1)***	2.18 (0.9)*
Unknown	0.74 (0.3)**	0.51 (0.2)*	4.33 (1.6)**	2.06 (1.3)	0.70 (0.3)*	0.58 (0.2)*	3.69 (1.8)*	3.61 (1.4)*
Wks*Biweekly to monthly	0.00 (0.0)	0.00 (0.0)	0.08 (0.0)	0.08 (0.0)	-0.01 (0.0)	-0.01 (0.0)	-0.06 (0.1)	-0.06 (0.1)
Wks*Weekly to biweekly	-0.01 (0.0)	-0.01 (0.0)	0.04 (0.1)	0.04 (0.1)	-0.01 (0.0)	-0.01 (0.0)	-0.06 (0.1)	-0.06 (0.1)
Wks*More than weekly	-0.05 (0.0)***	-0.05 (0.0)***	-0.16 (0.1)*	-0.16 (0.1)*	-0.03 (0.0)*	-0.03 (0.0)*	-0.16 (0.1)*	-0.16 (0.1)*
Other CT provider visits								
Monthly to < monthly	0.37 (0.1)**	0.22 (0.1)*	0.61 (0.8)	0.16 (0.6)	0.40 (0.1)**	0.23 (0.1)*	1.46 (0.8)	0.55 (0.7)
Biweekly up to monthly	0.42 (0.2)*	0.19 (0.1)	2.21 (1.1)*	1.43 (0.8)	0.33 (0.2)	0.10 (0.1)	0.21 (1.2)	-1.23 (0.9)
> Weekly up to biweekly	0.64 (0.2)***	0.33 (0.1)*	5.66 (1.0)***	4.43 (0.8)***	0.70 (0.2)***	0.45 (0.1)**	4.31 (1.1)***	2.89 (0.9)***
Unknown	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)
Wks* Monthly to < monthly	0.00 (0.0)	0.00 (0.0)	0.04 (0.0)	0.04 (0.0)	-0.01 (0.0)	-0.01 (0.0)	-0.11 (0.1)*	-0.11 (0.1)*
Wks* Biweekly to monthly	-0.01 (0.0)	-0.01 (0.0)	0.00 (0.1)	0.00 (0.1)	0.01 (0.0)	0.01 (0.0)	0.13 (0.1)	0.14 (0.1)
Wks*>Weekly to biweekly	0.01 (0.0)	0.01 (0.0)	0.12 (0.1)	0.12 (0.1)	0.00 (0.0)	0.00 (0.0)	0.11 (0.1)	0.11 (0.1)
Medical provider visit								
Monthly to < monthly	0.58 (0.1)***	0.20 (0.1)	4.75 (0.8)***	1.78 (0.6)**	0.35 (0.1)*	-0.06 (0.1)	4.20 (0.9)***	1.17 (0.7)
Biweekly up to monthly	1.46 (0.3)***	0.78 (0.2)***	15.40 (1.7)***	8.36 (1.3)***	1.41 (0.3)***	0.73 (0.3)**	17.24 (2.0)***	10.02 (1.6)***
> Weekly up to biweekly	1.09 (0.5)*	0.35 (0.4)	13.10 (2.9)***	5.92 (2.3)**	0.30 (0.6)	-0.18 (0.5)	6.79 (3.9)	0.00 (3.1)
Unknown	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)	0.00 (0.0)
Wks* Monthly to < monthly	0.00 (0.0)	0.00 (0.0)	0.01 (0.1)	0.01 (0.1)	0.00 (0.0)	0.00 (0.0)	0.12 (0.1)*	0.12 (0.1)*
Wks* Biweekly to monthly	-0.02 (0.0)	-0.02 (0.0)	0.17 (0.1)	0.17 (0.1)	-0.01 (0.0)	-0.01 (0.0)	-0.08 (0.1)	-0.08 (0.1)
Wks*>Weekly to biweekly	-0.01 (0.0)	-0.02 (0.0)	0.07 (0.2)	0.06 (0.2)	0.10 (0.1)	0.10 (0.1)	0.99 (0.3)***	0.99 (0.3)***
Years of Pain (Ref: < 1 Year)								
1 to < 2 Years	-	0.11 (0.2)	-	-0.73 (1.1)	-	0.04 (0.2)	-	1.64 (1.3)
2 to < 5 Years	-	0.04 (0.1)	-	-0.69 (0.9)	-	0.16 (0.2)	-	2.51 (1.0)*
5 to < 10 Years	-	0.02 (0.1)	-	-1.37 (0.9)	-	0.19 (0.2)	-	2.02 (1.0)*

Appendix Table 1. Results of hierarchical linear models for each outcome: models adding the impact of frequency of other types of visits and other patient characteristics now showing the estimated coefficients for other patient characteristics not included in Table 3. (continued)

	LBP Rating (n = 1361)		ODI (n = 1362)		NP Rating (n = 1214)		NDI (n = 1213)	
	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables
10+ Years	-	0.10 (0.1)	-	-0.07 (0.8)	-	0.30 (0.1)*	-	3.30 (0.9)***
Unknown	-	0.15 (0.2)	-	-0.01 (1.6)	-	0.05 (0.3)	-	3.65 (1.7)*
Both back & neck pain	-	0.14 (0.1)	-	-0.05 (0.5)	-	0.15 (0.1)	-	0.26 (0.6)
-% time doing heavy labor (Ref: None)								
-Non-work: > 0% to < 20%	-	0.18 (0.1)	-	1.40 (0.7)	-	0.08 (0.1)	-	2.00 (0.8)*
Workplace: > 0% to < 20%	-	0.14 (0.1)	-	-0.69 (0.6)	-	0.09 (0.1)	-	1.17 (0.7)
Non-work: > 20%	-	0.21 (0.2)	-	0.20 (1.3)	-	0.37 (0.2)	-	2.24 (1.6)
Workplace: > 20%	-	0.34 (0.1)**	-	-0.68 (0.8)	-	0.41 (0.1)**	-	1.33 (0.9)
Missing	-	0.05 (0.1)	-	1.66 (0.9)	-	0.13 (0.1)	-	2.30 (1.0)*
Surgery unsuccessful	-	0.49 (0.6)	-	9.92 (3.6)**	-	-1.07 (0.7)	-	1.40 (4.5)
Surgery unknown	-	-0.25 (0.3)	-	0.25 (2.1)	-	0.59 (0.5)	-	2.73 (3.1)
Often/always medication use past 6 months:								
OTC pain medications	-	0.06 (0.1)	-	0.41 (0.5)	-	-0.09 (0.1)	-	0.61 (0.5)
Rx pain medications	-	-0.03 (0.1)	-	-0.30 (1.0)	-	-0.05 (0.2)	-	0.09 (1.1)
Narcotic medications	-	0.30 (0.2)	-	3.44 (1.2)**	-	0.38 (0.2)	-	-0.24 (1.4)
Often/always exercise past 6 months	-	0.00 (0.1)	-	-1.45 (0.5)**	-	0.21 (0.1)*	-	-0.30 (0.5)
New patient (< 30 days)	-	0.06 (0.1)	-	0.30 (0.8)	-	0.34 (0.1)*	-	1.73 (0.9)
Unknown time w/chiro	-	-0.08 (0.1)	-	-1.10 (0.6)	-	0.11 (0.1)	-	-0.54 (0.7)
Ended care during 3 mos	-	-0.31 (0.1)*	-	-1.82 (0.9)*	-	-0.26 (0.2)	-	0.71 (1.2)
Believe pain is chronic	-	0.24 (0.1)***	-	1.85 (0.5)***	-	0.19 (0.1)*	-	1.83 (0.5)***
Pain w/o chiropractic	-	0.31 (0.0)***	-	1.17 (0.1)***	-	0.37 (0.0)***	-	1.51 (0.1)***
Exercise unsafe: agree to strongly agree	-	-0.04 (0.2)	-	3.00 (1.0)**	-	-0.26 (0.2)	-	2.05 (1.2)
Pain self-efficacy (1-10)	-	-0.05 (0.0)*	-	-1.57 (0.2)***	-	-0.02 (0.0)	-	-1.26 (0.2)***
Expect care successful	-	-0.43 (0.1)***	-	-0.84 (0.6)	-	-0.28 (0.1)**	-	-0.01 (0.7)
Expect a lot+ improvement	-	-0.12 (0.1)	-	-0.96 (0.5)	-	-0.02 (0.1)	-	-0.72 (0.6)
Pain worry mod to all time	-	0.12 (0.1)	-	0.42 (0.8)	-	-0.15 (0.1)	-	1.31 (0.9)
Depression (PROMIS)	-	0.00 (0.1)	-	0.86 (0.6)	-	-0.03 (0.1)	-	1.32 (0.7)*
Affective distress (0-7)	-	0.13 (0.0)***	-	0.78 (0.2)**	-	0.21 (0.0)***	-	1.16 (0.3)***
Catastrophizing (0-12)	-	0.12 (0.0)***	-	0.77 (0.1)***	-	0.17 (0.0)***	-	0.98 (0.2)***



## Chiropractic Visit Frequency Impact on Outcomes

Appendix Table 1. Results of hierarchical linear models for each outcome: models adding the impact of frequency of other types of visits and other patient characteristics now showing the estimated coefficients for other patient characteristics not included in Table 3. (continued)

	LBP Rating (n = 1361)		ODI (n = 1362)		NP Rating (n = 1214)		NDI (n = 1213)	
	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables	Adding other visit frequency	Adding all other variables
Average age in years	-	0.00 (0.0)	-	0.13 (0.0)***	-	-0.01 (0.0)**	-	0.05 (0.0)*
Gender: Female	-	0.21 (0.1)**	-	1.31 (0.5)*	-	0.22 (0.1)*	-	3.14 (0.6)***
Education: < 4-yr degree	-	-0.15 (0.1)*	-	-0.95 (0.5)*	-	-0.04 (0.1)	-	0.07 (0.5)
Partitioned Variance								
Clinic (Intercept)	0.02 (0.0)	0.01 (0.0)	5.48 (2.1)	1.84 (1.1)	0.07 (0.0)	0.00 (0.0)	3.80 (2.3)	0.57 (1.3)
ID (Weeks)	0.01 (0.0)	0.01 (0.0)	0.21 (0.0)	0.21 (0.0)	0.01 (0.0)	0.01 (0.0)	0.23 (0.0)	0.24 (0.0)
ID (Intercept)	2.23 (0.1)	1.02 (0.1)	99.93 (4.6)	52.38 (2.7)	2.64 (0.1)	1.13 (0.1)	111.07 (5.5)	58.86 (3.4)
Residual	1.61 (0.0)	1.61 (0.0)	27.76 (0.5)	27.73 (0.5)	1.60 (0.0)	1.60 (0.0)	32.65 (0.7)	32.62 (0.7)
D (df) for main effects	45.9 (8) ***	300.7 (33) ***	92.1 (8) ***	402.0 (33) ***	25.6 (8) ***	288.4 (33) ***	64.5 (8) ***	355.9 (33) ***
D (df) for interactions	0.8 (6)	-61.5 (32)	3.6 (6)	-25.4 (32)	2.5 (6)	-54.4 (32)	15.3 (6) ***	-12.2 (32)

BL = baseline; CT = complementary therapy providers—here, most often massage therapists; D = deviance statistic with its degrees of freedom (number of parameters added over previous model), distributed  $\chi^2$ ; LBP = low back pain, 0-10 scale; NDI = Neck Disability Index, 0-100 scale; NP = neck pain, 0-10 scale; ODI = Oswestry Disability Index, 0-100 scale; OTC = over-the-counter medications; Rx = prescription medications

\*The estimated coefficient or deviance statistic is statistically significant at  $P < 0.05$ .

\*\*The estimated coefficient or deviance statistic is statistically significant at  $P < 0.01$ .

\*\*\*The estimated coefficient or deviance statistic is statistically significant at  $P < 0.001$ .