

Opinion Survey

Training and Credentialing Standards for Minimally Invasive Spinal Surgery Techniques: Results of a Survey

Kai-Uwe Lewandrowski, MD¹, João Paulo Bergamaschi, MD², Albert E. Telfeian, MD, PhD³, Paulo Sérgio Teixeira de Carvalho, MD, PhD⁴, and Jorge Felipe Ramírez León, MD⁵

From: ¹Center for Advanced Spine Care of Southern Arizona and Surgical Institute of Tucson; ²Atualli Spine Care-Sao Paulo/Brazil, Alameda Santos, São Paulo-SP, Brazil; ³Department of Neurosurgery, Rhode Island Hospital, The Warren Alpert Medical School of Brown University, Providence, RI; ⁴Pain and Spine Minimally Invasive Surgery Service at Gaffre e Guinle University Hospital, Rio de Janeiro, Brazil; ⁵Minimally Invasive Spine Center, Bogotá, D.C., Colombia, Reina Sofia Clinic, Bogotá, D.C., Colombia

Address Correspondence:
Kai-Uwe Lewandrowski, MD
Center for Advanced Spine Care
of Southern Arizona
4787 E. Camp Lowell Dr.
Tucson, AZ 85712
E-mail:
business@tucsonspine.com

Disclaimer: The views expressed in this article represent those of the authors and no other entity or organization. The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. There was no external funding in the preparation of this manuscript.

Conflict of interest: Each author certifies that he or she, or a member of his or her immediate family, has no commercial association (i.e., consultancies, stock ownership, equity interest, patent/licensing arrangements, etc.) that might pose a conflict of interest in connection with the submitted manuscript.

Manuscript received: 07-02-2022

Background: The controversy continues on how to best become proficient in contemporary minimally invasive spinal surgery techniques (MISST). Postgraduate training programs typically lag behind the innovation. Other subspecialty spine care providers often compete with spine surgeons particularly when they do not offer the treatments needed by their patients. The public debate centers around who should be taught and credentialed in providing surgical spine care.

Objectives: The purpose of this study was to conduct an opinion survey amongst spine care providers regarding the learning curve of MISST and which credentialing standards should be established.

Setting: Surgeon online opinion survey sent by email, and chat groups in social media networks, including WeChat, WhatsApp, and LinkedIn.

Methods: Surgeons were asked the following questions: 1) Do you think MISS is harder to learn compared to open surgery? 2) Do you perform MISS? 3) What type of MISS do you perform? 4) If you perform endoscopic surgery, which approach(es)/technique(s) do you employ? 5) In your opinion, where does the innovation take place? 6) Where should MISST be taught? 7) Do you think mastering the MISST learning curve and surgeon skill level affect patient outcomes? 8) Which credentialing criteria do you recommend? Demographic data of responding surgeons, including age, postgraduate training and years from graduation, and practice setting, were also obtained. Descriptive statistics were employed to count the responses and compared to the surgeon's training using statistical package SPSS Version 27.0 (IBM Corporation, Armonk, NY).

Results: The online survey was viewed by 806 surgeons, started by 487, and completed by 272, yielding a completion rate of 55.9%. Orthopedic surgeons comprised 52.6% (143/272) of respondents, followed by 46.7% (127/272) neurosurgeons, and 0.7% pain management physicians (2/272). On average, respondents had graduated from a postgraduate training program 15.43 ± 10.13 years. Nearly all respondents employed MISST (252/272; 92.8%) and thought that proficiency in MISST affects patients' outcomes (270/272; 98.2%). Some 54.1% (146/270) opined that MISS is more challenging to learn than traditional open spine surgery. Preferred credentialing criteria were 1) number of MISST cases (87.5%; 238/272), b) skill level (69.9%; 190/272), and c) proficiency assessment (59.9%; 163/272). A case log review (42.3%; 116/272) or an oral examination (26.1%; 71/272) was not favored by surgeons. Surgeons reported academia (43.4%; 116/267) and private practice (41.2%; 110/267) as the centers of innovation. Only 15.4% (41/267) of respondents opined that industry was the main driver over innovation.

Limitations: Geographical and cultural biases may impact the opinions of responding surgeons.

Conclusions: Respondents preferred case volume, skill level, and proficiency assessment as credentialing criteria. Surgeons expect academic university programs and specialty societies to provide the necessary training in novel MISST while working with governing boards to update the certification programs.

Revised manuscript received:
10-10-2022
Accepted for publication:
11-10-2022

Free full manuscript:
www.painphysicianjournal.com

Key words: Surgeon opinion survey, minimally invasive spinal surgery, learning curve, credentialing, teaching standards

Pain Physician 2023: 26:29-37

As the demand for spine care rises (1-3), the competition between spine surgeons, which include nonsurgeons, such as anesthesiologists, physical therapists, physiatrists, Interventional radiologists, and new competing specialties, who provide interventional and procedural pain management treatments, intensifies (4-7). By default, these treatments, some of them consisting of innovative minimally invasive spinal surgery techniques (MISST), are often left to nonsurgeons, as traditionally trained spine surgeons usually do not offer alternatives to extensive and aggressive operations to patients (8-10). The debate on who should be providing what type of spine care typically heats up when surgeons are asked to take care of a complication or inferior clinical outcome that a nonsurgeon could not revise (11). Training standards and credentialing requirements are vehemently debated when other subspecialties invade the "surgical turf" of spine surgeons and when referrers send their patients to alternate providers. Examples of such trends include the rapid growth of spinal cord stimulators and pain pumps (12). Many patients who ordinarily would have been referred for surgical decompression, some of which would face the additional prospect of a more complex fusion, are often referred for such procedures. Nowadays, some of them are done performed by accomplished MI surgeons, but all new competitors for patients with or without formal training are now in the mix. Sacroiliac joint fusions are also increasingly performed by surgeons, as well as pain management physicians, who now use surgical codes to be reimbursed for the procedure (13-18). Another example is spinal endoscopy (19,20). The percutaneous nature of the surgical approach through a < 1-centimeter incision may be tempting to some interventionalists to expand their procedural portfolio (21). Some vendors have a product kit lineup ranging from nonvisualized combination decompression radiofrequency procedure done under fluoroscopic control to visualized videoendoscopic decompression, which by most would be considered surgery and not an interventional procedure. In some instances, the product components are even identical between these applications (22,23). In other words, the line between interventional spine

care and spinal surgery is becoming increasingly fuzzy, creating the need to redefine training and credentialing standards to catch up with this fast-moving field. Most of the time, local medical staff offices are left to define their own standards when credentialing a provider for innovative procedures or surgeries with which they have no experience or prior track record.

The overlap in treating the same painful degenerative spine conditions between subspecialties creates controversy and the need to better understand the issues at stake that drive the innovation dynamic. Traditionally, governing boards and scientific surgeon organizations have defined the curriculum requirements before an applicant is even admitted to board certification examination (24). After completing the required postgraduate training, becoming board certified in a surgical subspecialty is no small undertaking and requires ongoing life-long learning via minimum continued medical education requirements. The latter is typically a process of several consecutive steps that have to be taken between oral and written tests. Over the professional lifetime, surgeons might recertify several times to remain in good standing with the governing board that certifies to their ability to treat members of the public effectively and safely while adhering to the highest professional standards. It is, therefore, easy to understand that any disregard of such hard-earned certification requirements by others may be looked upon critically and potentially considered inappropriate. In some cases, nonsurgeons were sued by their patients and have lost their medical licenses for practicing outside the purview of their training and board certification. The purpose of this opinion survey study relevant to implementation and quality control with modern MISST was to get an up-to-date snapshot in time on the opinions of those who are at the center of credentialing and learning debate-spine surgeons.

METHODS

The authors solicited responses to an online survey via email and chat groups in social networks, including Facebook, WeChat, WhatsApp, and LinkedIn. The survey was available online and distributed via a link distributed through these social network media. Upon

clicking on the link, the prospective respondent was taken to the typeform Web site at www.typeform.com. The typeform service works on any computer, laptop, and any handheld devices, such as an iPad, or a cellular smartphone with multiple Web browser user-interface platforms. The questions were aimed at soliciting information deemed important to the understanding of current practice patterns and opinions regarding innovative MISST, training, and credentialing; whereas, another question requested demographic information of the respondent regarding postgraduate residency and fellowship training, and their practice setting. The survey was constructed of either simple "YES" or "NO" questions or simple multiple-choice questions, some of which with multiple possible answers. Surgeons were asked the following questions:

1. Do you think MISS is harder to learn compared to open surgery?
YES/NO
2. Do you perform MISS?
YES/NO
3. What type of MISS do you perform?
 - i. Tubular retractor system
 - ii. Mini-open surgery
 - iii. Endoscopic surgery
 - iv. Industry
 - v. Other
4. If you perform endoscopic surgery, which approach(es)/technique(s) do you employ?
 - i. Transforaminal
 - ii. Interlaminar
 - iii. Full endoscopic
 - iv. Over the top
 - v. Unilateral biportal
 - vi. Other
5. In your opinion, where does the innovation take place?
 - i. Academia
 - ii. Private practice
 - iii. Other
6. Where should MISST be taught?
 - i. University program
 - ii. Private practice
 - iii. Specialty society
 - iv. Industry
 - v. Other
7. Do you think mastering the MISST learning curve and surgeon skill level affect patient outcomes?
YES/NO
8. Which credentialing criteria do you recommend?
 - i. Number of cases
 - ii. Proficiency assessment
 - iii. Oral examination
 - iv. Case log review
 - v. Skill assessment
 - vi. Other
9. What is your training?
 - i. Orthopedic spine surgeon
 - ii. Neurosurgeon
 - iii. Other
10. What is your age?
11. How many years ago did you graduate from a post-graduate training program (residency/fellowship)?
12. Tell us about your practice setting:
 - i. Hospital employed
 - ii. University employed
 - iii. Private practice
 - iv. Private practice with academic appointment
 - v. Other
13. What's your country of residence?

The double-blinded survey ran from October 8 to December 2, 2021. Upon completion of the survey, the responses were downloaded in an Excel file format and imported into SPSS Version 27.0 (IBM Corporation, Armonk, NY) statistical software package for further data analysis. Descriptive statistics included counts, means, range, and standard deviation as well as percentages. Missing answers were not included in the analysis. Wherever applicable, a *P* value of 0.05 or less was considered statistically significant, and a confidence interval of 95% was considered for all statistical tests.

RESULTS


The online survey was viewed by 806 respondents, started by 487, and completed by 272. The completion rate was 55.9% and the average time to complete the survey was 04 minutes and 57 seconds. Just over


half of surgeons (143/272; 52.6%) who participating in this survey were orthopedic surgeons, followed by neurosurgeons (127/272; 46.7%). Two respondents were pain management physicians (2/272; 0.7%). These respondents were between the ages of 35-44 years; 41.5%, and between the ages of 45-55 years; 33.1%, at the time. The remaining age groups were represented as follows: 55-64 years; 14%, 25-34 years; 4.4%, 65-74 years; 4.4%, and 18-24 years; 0.7%. On average, respondents had graduated from a postgraduate training program 15.43 ± 10.13 ranging from 1-49 years. In decreasing order of relevance, surgeons reported from Brazil (71/272; 26.7%), China (58/272; 21.8%), Mexico (33/272; 12.4%), South Korea (20/272; 7.5%), Chile (11/272; 4.1%), Paraguay (10/272; 3.8%), India (9/272; 3.4%), United States (8/272; 3%), Argentina (7/272; 2.6%), Colombia (7/272; 2.6%), Bolivia (6/272; 2.3%), El Salvador (3/272; 1.1%), Spain (3/272; 1.1%), Bangladesh (2/272; 0.8%), Malaysia (2/272; 0.8%), Netherlands (2/272; 0.8%), Portugal (2/272; 0.8%), Thailand (2/272; 0.8%), Algeria (1/272; 0.4%), Costa Rica (1/272; 0.4%), Dominican Republic (1/272; 0.4%), Indonesia (1/272; 0.4%), Nepal (1/272; 0.4%), Philippines (1/272; 0.4%), Singapore (1/272; 0.4%), Syria (1/272; 0.4%), United Kingdom (1/272; 0.4%), and Vietnam (1/272; 0.4%). The majority (108/272; 39.7%) of these responding surgeons were self-employed and worked in a private practice setting. The second largest group (69/272; 25.4%) was comprised of hospital

employed surgeons. Forty- Forty-five (16.2%) were private practice surgeons with an academic appointment, and another 44 surgeons were university employed. Only 6 surgeons (2.2%) were employed by a large physician-owned group (Fig. 1).

The majority of respondents performed MISS (252/272; 92.8%), and nearly all respondents (270/272; 98.2%) opined that proficiency in MISST affects patients' outcomes. However, only 54.1% (146/270) opined that MISS is more challenging to learn than traditional open spine surgery (124/270; 45.9%). Considering that multiple answers were allowed, 73.5% (200/272) of respondents performed endoscopic spine surgery, 57.7% (157/263) performed mini-open surgery, and 53.3% (145/263) employed a tubular retractor system during their spine surgeries. A breakdown of the practiced endoscopic spine surgeries revealed that 56.1% (145/263) of responding physicians performed interlaminar endoscopy, 53.2% (140/263) transforaminal endoscopy, 45.2% (119/263) full endoscopy, 23.6% (62/263) over the top, and 18.6% (49/263) unilateral biportal endoscopy. Only 15.2% (40/263) did not perform endoscopic spine surgery (Fig. 2).

When asked where innovation took place, 43.4% (116/267) thought that new inventions and clinical protocols emerged primarily in academia; whereas, 41.2% (110/267) considered private practice the primary place where clinical innovation in spine surgery played out. Only 15.4% (41/267) respondents opined that industry





was the main driver over innovation. When it came to who should orchestrate the teaching of modern MISST, private practice was even less relevant to survey respondents. Only 23.9% of respondents (65/272) felt that private practice was the most appropriate setting for innovation. The 2 settings for teaching novel MISST desired by survey respondents the most were a specialty society (35.7%; 97/272) and a university program (35.7%; 97/272). Only 4.8% (13/272) of them preferred that the industry orchestrate the training. The credentialing criteria were at the heart of this survey. The majority (87.5%; 238/272) of surgeons thought

that the most critical credentialing criterion was the number of cases performed by a surgeon. Skill (69.9%; 190/272) and proficiency assessment (59.9%; 163/272) were other preferred credentialing criteria. A case log review (42.3%; 116/272) or an oral examination (26.1%; 71/272) did not find majority support by surgeons (Fig. 2).

DISCUSSION

Findings of this opinion survey of 272 responding surgeons who returned completed an online questionnaire indicate that spine surgeons did find MISST

somewhat more challenging to learn and that skill level impacts clinical outcomes, thereby substantiating the authors' call for credentialing standards. Surgeons overwhelmingly voted for using the number of MISST cases (87.5%; 238/272) as the primary criterion for credentialing spine surgeons in novel MISST. The respondents also supported skill level (69.9%; 190/272) and proficiency assessment (59.9%; 163/272). A case log review (42.3%; 116/272) or an oral examination (26.1%; 71/272) did not find majority support by surgeons. The significance of these findings is that postgraduate and society specialty training programs should structure their curriculum to respond to surgeons' demand of

high caseload and skill-building activities to improve proficiency. Coincidentally, spine surgeons also identified academia and a specialty society as the preferred venues where the training should occur, and the credentialing be administered. Spine surgeons did not accept MISST training by industry (Fig. 3).

The online survey proved an effective means of collecting tangible clinical information by reaching 806 surgeons, of which 487 started and 272 completed the survey (Fig. 1). This methodology has been employed by the authors in several studies (25-30) that have been published in peer-reviewed journals. It was instrumental in gathering large amounts of data quickly across




Fig. 3. The percentage breakdown of responses to survey questions on MISST practice patterns and relevance of credentialing and training standards with multiple possible answers given by the 272 participating spine surgeons is shown. a) The majority of responding surgeons preferred case volume (87.5%; 238/272), skill level (69.9%; 190/272), and proficiency assessment (59.9%; 163/272) as credentialing criteria. b) Most surgeons worked in private practice (108/272; 39.7%), followed by hospital employment (69/272; 25.4%), and private practice with an academic appointment (45/272; 16.2%). A sizeable physician-owned group employed only 6 surgeons (2.2%). c) Respondents were made up of orthopedic surgeons (52.6%; 143/272), followed by neurosurgeons (46.7%; 127/272). Two respondents were pain management physicians (0.7%; 2/272). MISST, minimally invasive spinal surgery.

many cultural and geographical barriers, particularly when assessing the most appropriate management of rare complications (31) or understanding surgeons' and health care facility needs when establishing an endoscopic spinal surgery program (29). One of the authors' study assessed the incidence of durotomies based on a total of 64,470 lumbar endoscopies (25). This type of "big data" analysis is typically impossible at a single center site and may even prove difficult in multicenter trials. The present survey reached physicians working in hundreds of health care facilities worldwide.

The high completion rate of 55.9% is much higher than response rates reported with traditional in-mail surveys—5% or less (32). The literature shows that response rates may vary. The average response rate for an in-person survey is 57%, a mail survey at 50%, an email survey at 30%, an online survey at 29%, a telephone survey at 18%, and an in-app survey at 13%, with an overall average survey response rate of 33% (32-35). A survey mailed to 2,048 patients had a response rate of 46% (36). Web-based surveys have achieved a response rate of 60% (32). Paper surveys have a response rate of 49.5% (37). An online patient satisfaction survey of 9,975 Medicare beneficiaries with 434 unduplicated survey submissions had a response rate of 4.3% (36). Clinical opinion survey research amongst spine surgeons showed response rates ranging from 49% (n = 51) obtained in a study on surgical management of spinal stenosis amongst Norwegian spine surgeons (38) to 61.3% (n = 357) amongst members of the Scoliosis Research Society surveyed on use of MISST in the treatment of adult spinal deformity (39). In comparison, another sizeable online survey study (40) sent to trauma, spine, and craniomaxillofacial surgeon members of the Association of Osteosynthesis Foundation, yielded 1,212 valid and completed submissions at an overall response rate of 4.1%. These examples illustrate that there can be a wide range of survey response rates and that 55.9% obtained in this study is higher than the average reported for clinical online survey research. Research also shows that higher response rates do not necessarily translate into higher survey accuracy since missing answers do not occur randomly (35,41).

This study likely suffered from some bias. The authors did their best to minimize it. Intuition and hindsight bias is common amongst the respondents and investigators with this type of data acquisition (42,43). The authors captured opinions almost exclusively from orthopedic spine surgeons (52.6%) and neurosurgeons

(46.7%). Only 2 pain management physicians (0.7%) entered data into the survey. Several explanations for the low percentage of pain physicians responding to this survey seem plausible: 1) There are likely fewer pain physicians performing endoscopic surgery than orthopedic and neurosurgeons. 2) The authors are all surgeons, and their selection bias could have preferentially distributed the survey to other surgeons. Ultimately, another survey directly addressed to pain management physicians could provide more conclusive insights into whether there is any difference regarding endoscopic spine surgery credentialing criteria between spine care providers with different postgraduate training backgrounds. Responses were blinded, and the investigators of this study had no way of anticipating any causes for regional variations in physician preferences regarding endoscopic spinal surgery credentialing criteria. Understanding the incoming data in real-time was essential to the authors since surgeons from some countries (Brazil; 26.7%, China; 21.8%, Mexico; 12.4%, South Korea; 7.5%, Chile; 4.1%, Paraguay; 3.8%, India; 3.4%, United States; 3%), were somewhat overrepresented in the survey making up 82.7% of all respondents. As recently shown, these nations are also the countries where spinal endoscopy is most accepted and considered mainstream (27). Besides, the effect size of agreements or disagreements was unknown when launching the survey. The rapid response dynamic observed by the authors (Fig. 1) suggests the study findings are, in fact, representative of current opinions regarding endoscopic spinal surgery criteria amongst physicians involved in the management of spine pain the world over.

CONCLUSIONS

This online survey reached 272 spine care providers in just 2 months, proving that surgeon opinions can be readily solicited on hand-held devices. Surgeons preferred case volume, skill level, and proficiency assessment as credentialing criteria. These criteria would virtually guarantee that surgical procedures are done by well-trained surgeons who know how to handle perioperative problems, complications, and perform revision surgeries if and when needed. Surgeons also indicated that they expect academic university programs and specialty societies to provide the necessary training and work hand-in-hand with local, state, and federal governing boards to update the certification programs to keep up with the fast-moving innovative field of MISS.

REFERENCES

1. Backstrom KM, Whitman JM, Flynn TW. Lumbar spinal stenosis-diagnosis and management of the aging spine. *Man Ther* 2011; 16:308-317.
2. Graves JM, Fulton-Kehoe D, Martin DP, et al. Factors associated with early magnetic resonance imaging utilization for acute occupational low back pain: A population-based study from Washington State workers' compensation. *Spine (Phila Pa 1976)* 2012; 37:1708-1718.
3. Haldeman S, Johnson CD, Chou R, et al. The Global Spine Care Initiative: Care pathway for people with spine-related concerns. *Eur Spine J* 2018; 27:901-914.
4. Arts MP, Peul WC, Koes BW, et al. Management of sciatica due to lumbar disc herniation in the Netherlands: A survey among spine surgeons. *J Neurosurg Spine* 2008; 9:32-39.
5. Atlas SJ, Deyo RA, Keller RB, et al. The Maine Lumbar Spine Study, Part II. 1-year outcomes of surgical and nonsurgical management of sciatica. *Spine (Phila Pa 1976)* 1996; 21:1777-1786.
6. Atlas SJ, Keller RB, Chang Y, et al. Surgical and nonsurgical management of sciatica secondary to a lumbar disc herniation: Five-Year outcomes from the Maine Lumbar Spine Study. *Spine (Phila Pa 1976)* 2001; 26:1179-1187.
7. Atlas SJ, Keller RB, Wu YA, et al. Long-Term outcomes of surgical and nonsurgical management of lumbar spinal stenosis: 8 to 10 year results from the Maine Lumbar Spine Study. *Spine (Phila Pa 1976)* 2005; 30:936-943.
8. Lurie JD, Berven SH, Gibson-Chambers J, et al. Patient preferences and expectations for care: Determinants in patients with lumbar intervertebral disc herniation. *Spine (Phila Pa 1976)* 2008; 33:2663-2668.
9. Lurie JD, Henderson ER, McDonough CM, et al. Effect of expectations on treatment outcome for lumbar intervertebral disc herniation. *Spine (Phila Pa 1976)* 2016; 41:803-809.
10. Menendez JY, Omar NB, Chagoya G, et al. Patient satisfaction in spine surgery: A systematic review of the literature. *Asian Spine J* 2019; 13:1047-1057.
11. Whitmore RG, Stephen J, Stein SC, et al. Patient comorbidities and complications after spinal surgery: A societal-based cost analysis. *Spine (Phila Pa 1976)* 2012; 37:1065-1071.
12. Atallah J, Armah FA, Wong D, et al. Use of spinal cord stimulator for treatment of lumbar radiculopathy in a patient with severe kyphoscoliosis. *Pain Physician* 2008; 11:555-559.
13. Boswell MV, Shah RV, Everett CR, et al. Interventional techniques in the management of chronic spinal pain: Evidence-Based practice guidelines. *Pain Physician* 2005; 8:1-47.
14. Cognetti DJ, Jorgensen AY. Minimally invasive sacroiliac fusion revision: A technique guide. *Int J Spine Surg* 2021; 15:274-279.
15. Lee DW, Patterson DG, Sayed D. Review of current evidence for minimally invasive posterior sacroiliac joint fusion. *Int J Spine Surg* 2021; 15:514-524.
16. Lorio M, Kube R, Araghi A. International Society for the Advancement of Spine Surgery Policy 2020 update-minimally invasive surgical sacroiliac joint fusion (for chronic sacroiliac joint pain): Coverage indications, limitations, and medical necessity. *Int J Spine Surg* 2020; 14:860-895.
17. Lorio MP. Editor's introduction: Update on current sacroiliac joint fusion procedures: Implications for appropriate current procedural terminology medical coding. *Int J Spine Surg* 2020; 14:853-859.
18. Polly DW Jr, Holton KJ. Minimally invasive sacroiliac joint fusion: A lateral approach using triangular titanium implants and navigation. *JBJs Essent Surg Tech* 2020; 10:e19.00067.
19. Jasper GP, Francisco GM, Telfeian AE. A retrospective evaluation of the clinical success of transforaminal endoscopic discectomy with foraminotomy in geriatric patients. *Pain Physician* 2013; 16:225-229.
20. Jasper GP, Francisco GM, Telfeian AE. Transforaminal endoscopic discectomy with foraminoplasty for the treatment of spondylolisthesis. *Pain Physician* 2014; 17:E703-E708.
21. Chung AS, Kimball J, Min E, et al. Endoscopic spine surgery-increasing usage and prominence in mainstream spine surgery and spine societies. *J Spine Surg* 2020; 6:S14-S18.
22. Lorio M, Kim C, Araghi A, et al. International Society for the Advancement of Spine Surgery Policy 2019-surgical treatment of lumbar disc herniation with radiculopathy. *Int J Spine Surg* 2020; 14:1-17.
23. Yeung A, Gore S. Endoscopically guided foraminal and dorsal rhizotomy for chronic axial back pain based on cadaver and endoscopically visualized anatomic study. *Int J Spine Surg* 2014; 8:23.
24. Lorio M, Clerk-Lamallice O, Beall DP, et al. International Society for the Advancement of Spine Surgery guideline-intraosseous ablation of the basivertebral nerve for the relief of chronic low back pain. *Int J Spine Surg* 2020; 14:18-25.
25. Lewandrowski KU, Hellinger S, De Carvalho PST, et al. Dural tears during lumbar spinal endoscopy: Surgeon skill, training, incidence, risk factors, and management. *Int J Spine Surg* 2021; 15:280-294.
26. Lewandrowski KU, Soriano-Sanchez JA, Zhang X, et al. Surgeon training and clinical implementation of spinal endoscopy in routine practice: Results of a global survey. *J Spine Surg* 2020; 6:S237-S248.
27. Lewandrowski KU, Soriano-Sanchez JA, Zhang X, et al. Regional variations in acceptance, and utilization of minimally invasive spinal surgery techniques among spine surgeons: Results of a global survey. *J Spine Surg* 2020; 6:S260-S274.
28. Lewandrowski KU, Soriano-Sanchez JA, Zhang X, et al. Surgeon motivation, and obstacles to the implementation of minimally invasive spinal surgery techniques. *J Spine Surg* 2020; 6:S249-S259.
29. Lewandrowski KU, Tieber F, Hellinger S, et al. Durability of endoscopes used during routine lumbar endoscopy: An analysis of use patterns, common failure modes, impact on patient care, and contingency plans. *Int J Spine Surg* 2021; 15:1147-1160.
30. Lewandrowski KU, Zhang X, Ramirez Leon JF, et al. Lumbar vacuum disc, vertical instability, standalone endoscopic interbody fusion, and other treatments: An opinion based survey among minimally invasive spinal surgeons. *J Spine Surg* 2020; 6:S165-S178.
31. Lewandrowski KU, Telfeian AE, Hellinger S, et al. Difficulties, challenges, and the learning curve of avoiding complications in lumbar endoscopic spine surgery. *Int J Spine Surg* 2021; 15:S21-S37.
32. Keller A. What is an acceptable survey response rate? *National Social Norms Center* 2014. <http://socialnorms.org/what-is-anacceptable-survey-response>

- rate
33. Keeter S, Hatley N, Kennedy C, Arnold Lau. What low response rates mean for telephone surveys. *Pew Research Center* 2017; 15:1-39.
 34. Ramshaw A. *The complete guide to acceptable survey response rates*. Accessed 04/01/2020. www.genroe.com/blog/acceptable-survey-response-rate-2/11504
 35. Sefferman A. *Mobile survey response rates: Averages and benchmarks by app category*. Apptentive 2016. www.apptentive.com/blog/mobile-survey-response-rates/
 36. Brandt NJ, Cooke CE, Sharma K, et al. Findings from a national survey of medicare beneficiary perspectives on the medicare part D medication therapy management standardized format. *J Manag Care Spec Pharm* 2019; 25:366-391.
 37. Odgaard L, Kothari M. Survey of oral nursing care attitudes, knowledge and practices in a neurorehabilitation setting. *J Oral Rehabil* 2019; 46:730-737.
 38. Weber C, Lønne G, Rao V, et al. Surgical management of lumbar spinal stenosis: A survey among Norwegian spine surgeons. *Acta Neurochir (Wien)* 2017; 159:191-197.
 39. Anand N, Agrawal A, Burger EL, et al. The prevalence of the use of MIS techniques in the treatment of adult spinal deformity (ASD) amongst members of the Scoliosis Research Society (SRS) in 2016. *Spine Deform* 2019; 7:319-324.
 40. Joeris A, Knoll C, Kalampoki V, et al. Patient-Reported outcome measurements in clinical routine of trauma, spine and craniomaxillofacial surgeons: Between expectations and reality: A survey among 1212 surgeons. *BMJ Open* 2018; 8:e020629.
 41. Mind-Blowing employee survey statistic. www.officevibe.com/blog/employee-surveys-infographic
 42. Henriksen K, Kaplan H. Hindsight bias, outcome knowledge and adaptive learning. *Qual Saf Health Care* 2003; 12(suppl 2):ii46-ii50.
 43. Zwaan L, Monteiro S, Sherbino J, et al. Is bias in the eye of the beholder? A vignette study to assess recognition of cognitive biases in clinical case workups. *BMJ Qual Saf* 2017; 26:104-110.

